



U.S. Department of Energy
Oakland Operations Office, Oakland, California

**FINAL
RADIONUCLIDE AIR EMISSION ANNUAL REPORT
(SUBPART H OF 40 CFR 61)
CALENDAR YEAR 2002**

for the

**LABORATORY FOR ENERGY-RELATED HEALTH RESEARCH
UNIVERSITY OF CALIFORNIA, DAVIS**

prepared for:

United States Department of Energy
NNSA Service Center
1301 Clay Street
Oakland, California 94612-5208

prepared by:

Weiss Associates
5801 Christie Avenue, Suite 600
Emeryville, California 94608-1827

April 16, 2003
Rev. 0

DOE Oakland Operations Contract DE-AC03-96SF20686

FINAL
RADIONUCLIDE AIR EMISSION ANNUAL REPORT
(SUBPART H OF 40 CFR 61)
CALENDAR YEAR 2002

for the
LABORATORY FOR ENERGY-RELATED HEALTH RESEARCH
UNIVERSITY OF CALIFORNIA, DAVIS

prepared for:

United States Department of Energy
NNSA Service Center
1301 Clay Street
Oakland, California 94612-5208

prepared by:

Weiss Associates
5801 Christie Avenue, Suite 600
Emeryville, California 94608-1827

April 16, 2003
Rev. 0

DOE Oakland Operations Contract DE-AC03-965F20686

Issued To: _____ Date: _____

Copy No.: _____ Controlled Uncontrolled

Approvals Page

FINAL
RADIONUCLIDE AIR EMISSION ANNUAL REPORT
(SUBPART H OF 40 CFR 61)
CALENDAR YEAR 2002

for the
LABORATORY FOR ENERGY-RELATED HEALTH RESEARCH
UNIVERSITY OF CALIFORNIA, DAVIS

prepared for:

U.S. Department of Energy
NNSA Service Center
1301 Clay Street
Oakland, California 94612-5208

prepared by:

Weiss Associates
5801 Christie Avenue, Suite 600
Emeryville, California 94608-1827

April 16, 2003
Rev. 0

Approved by: Robert O. Devany for Date: 4/16/03
Michael D. Dresen, C.E.G., C.H.G.
Program Manager
Weiss Associates

Approved by: Dolores D. Loll Date: 4/16/03
Dolores D. Loll, P.E.
Project Quality Assurance Manager
Weiss Associates

Approved by: Robert O. Devany Date: 4/16/03
Robert O. Devany, C.E.G., C.H.G.
Project Manager
Weiss Associates

CONTENTS

1.	FACILITY INFORMATION	1-1
1.1	Site Description	1-1
1.2	Source Description	1-3
1.2.1	Point Sources	1-3
1.2.2	Non-Point Sources	1-3
1.2.3	Western and Eastern Dog Pens Areas	1-3
1.2.4	Domestic Septic Systems 3 and 6	1-4
2.	AIR EMISSION DATA	2-1
2.1	Point Sources	2-1
2.2	Non-Point Sources	2-1
2.3	Particulate Matter-10	2-3
3.	DOSE ASSESSMENTS	3-1
3.1	Description of Dose Model	3-1
3.2	Summary of Input Parameters	3-2
4.	COMPLIANCE ASSESSMENT	4-1
4.1	Certification	4-1
5.	ADDITIONAL INFORMATION	5-1
6.	REFERENCES	6-1

TABLES

- Table 1. Summary of LEHR Site Potential Non-Point Diffuse Area Radionuclide Sources
- Table 2. Summary of Maximum Surface Soil Concentrations for Radionuclides Exceeding Background and PM₁₀ Emission Rates, LEHR Western Dog Pens Area
- Table 3. Summary of Maximum Surface Soil Concentrations for Radionuclides Exceeding Background and PM₁₀ Emission Rates, LEHR Eastern Dog Pens Area
- Table 4. Summary of Maximum Surface Soil Concentrations for Radionuclides Exceeding Background and PM₁₀ Emission Rates, LEHR Domestic Septic System 3
- Table 5. Summary of Maximum Surface Soil Concentrations for Radionuclides Exceeding Background and PM₁₀ Emission Rates, LEHR Domestic Septic System 6
- Table 6. Summary of On-Site Effective Dose Equivalent to Maximally Exposed Individual Resulting from Radionuclide Emissions from Each Potential Fugitive Dust Emission Non-Point Source
- Table 7. Summary of Estimated Collective Population Dose Resulting from Radionuclide Emissions from Each Fugitive Dust Emission Non-Point Source
- Table 8. CAP88-PC Inputs for the Western Dog Pens Area
- Table 9. CAP88-PC Inputs for the Eastern Dog Pens Area
- Table 10. CAP88-PC Inputs for the Domestic Septic System 3
- Table 11. CAP88-PC Inputs for the Domestic Septic System 6

FIGURES

Figure 1. LEHR Facility Location Map, UC Davis, California

Figure 2. LEHR Facility Potential and Known DOE Non-Point Source Areas

APPENDICES

Appendix A CAP88-PC Output Results

Appendix B DOE Supplemental Information

ACRONYMS AND ABBREVIATIONS

#/km ²	number per square kilometer
°C	degrees Celsius
°F	degrees Fahrenheit
Ac-228	actinium-228
AM	Air Monitoring Station
Am-241	americium-241
bgs	below ground surface
Bi-212	bismuth-212
Bi-214	bismuth-214
CAP88-PC	atmospheric dispersion/radiation dose calculation computer code (US EPA)
C-14	carbon-14
CERCLA	Comprehensive Environmental Response, Compensation and Liability Act
CFR	Code of Federal Regulations
Ci/yr	curies per year
cm/s	centimeters per second
cm/yr	centimeter(s) per year
Co-60	cobalt-60
Cs-137	cesium-137
cu yd	cubic yard(s)
DB	distribution box
DOE	United States Department of Energy
DSS	domestic septic system
DST	domestic septic tank
E	east
EDE	effective dose equivalent
EDPs	Eastern Dog Pens

Emis.	emission
ER/WM	Environmental Restoration/Waste Management
g/m ²	grams per square meter
g/m ² -sec	grams per square meter-seconds
hr	hour
H-3	tritium
ID	identification (number)
K-40	potassium-40
km	kilometers
km ²	square kilometer(s)
LEHR	Laboratory for Energy-Related Health Research
m	meter(s)
m/s	meters per second
m ²	square meter(s)
m ³	cubic meter (s)
Max.	maximum
mCi	milliCuries
MEI	maximally exposed individual
MOU	Memorandum of Understanding
mph	miles per hour
mrem/yr	millirem per year
mSv/yr	milliSievert per year
N	north
NNE	north - northeast
NNW	north - northwest
NESHAP	National Emissions Standards for Hazardous Air Pollutants
Pb-210	lead-210
Pb-212	lead-212
Pb-214	lead-214
pCi/g	picoCurie per gram

person-rem/yr	person - roentgen equivalent per year
PM ₁₀	particulate matter with an aerodynamic diameter less than or equal to 10 micrometers
Pu-241	plutonium-241
RA	removal action
Ra/Sr	radium/strontium
Ra-226	radium-226
S	south
Sr-90	strontium-90
Th-228	thorium-228
Th-234	thorium-234
Tl-208	thallium-208
U-233	uranium-233
U-234	uranium-234
U-235	uranium-235
U-238	uranium-238
UC Davis	University of California, Davis
UCL	upper confidence limit
US EPA	United States Environmental Protection Agency
W	west
WDPs	Western Dog Pens
µg	microgram(s)
µg/m ³	micrograms per cubic meter
µm	micrometer

US Department of Energy
Radionuclide Air Emission Annual Report
(Subpart H of 40 CFR 61)
Calendar Year 2002

Site Name: Laboratory for Energy-Related Health Research (LEHR)

Field Office Information

Office: NNSA Service Center
Address: 1301 Clay Street, Room 700 N
Oakland, CA 94612
Contact: Steve Black Phone: (510) 637-1595

Site Information

Operator: Weiss Associates
Address: University of California
Old Davis Road
One Shields Avenue
Davis, CA 95616-8615
Contractor Contact: Robert Devany Phone: (510) 450-6000
Project Manager
Weiss Associates
Site Contact: Kim Warren Phone: (530) 752-1348
Environmental Scientist
Weiss Associates

1. FACILITY INFORMATION

The Laboratory for Energy-Related Health Research (LEHR) is in compliance with the requirements of Title 40 Code of Federal Regulations (CFR) Part 61 Subpart H - National Emissions Standards for Hazardous Air Pollutants (NESHAP) for Emissions of Radionuclides from United States Department of Energy (DOE) Facilities. The NESHAP regulations require that radionuclide emissions not exceed levels that would result in an effective dose equivalent (EDE) of 10 millirem per year (mrem/yr).

This Radionuclide Air Emission Annual Report applies only to the DOE areas at the LEHR facility on the University of California, Davis (UC Davis) campus. For the purposes of this report, the DOE areas will be referred to as "the Site" and references to the LEHR include the entire 15-acre property where both DOE and UC Davis conduct activities. There are currently no point sources of radionuclide emission at the Site. Potential fugitive/area sources of radionuclide emissions were modeled using the United States Environmental Protection Agency (US EPA) atmospheric dispersion/radiation dose calculation computer code, CAP88-PC Version 1.0. Based on results from this model, the total estimated contribution to the maximum EDE from non-point source emissions for reporting year 2002 is estimated to be 3.8×10^{-4} mrem/yr (less than 0.01% of the 10 mrem/yr standard).

1.1 Site Description

The LEHR facility is located in Solano County, California, in the southeast quadrant of Section 21, Township 8 North, Range 2 East, Mount Diablo Base and Meridian. It is approximately 1.5 miles south of the town of Davis (Figure 1), and occupies about 15 acres on the southeast portion of the UC Davis campus. The LEHR facility is bounded by UC Davis research facilities, private farmland and the South Fork of Putah Creek. The southern boundary of the LEHR facility is the northern levee of the South Fork of Putah Creek.

The local climate is Mediterranean with mild winters and dry summers. Based on the site meteorological station data collected in 2002, the average winter (October through April) temperature was 51 degrees Fahrenheit (°F), and the average summer (May through September) temperature was 69°F. Precipitation data were also collected from the UC Davis climatological data center, located approximately one mile northwest of the LEHR facility. Based on the UC Davis data, the total precipitation for 2002 was 10.39 inches (with totals of 21.66, 16.07 and 10.53 inches recorded in 2001, 2000 and 1999, respectively.) Precipitation data from a local weather station for November and December of 2002 was not available. The sun shines approximately 95% of the time during daylight hours in the summer and about 45% of the time during daylight hours in the winter.

The prevailing wind direction is from the south, reflecting frequent incursions of marine air through the Carquinez Strait into the Sacramento Valley. Changes in wind direction are common, with winds from the northwest occurring diurnally. Several times a year, strong winds blow from the north, generally following the passage of Pacific storm systems (DOE, 1994a). The average 2002 wind speed recorded at the Site three-meter tower on the meteorological station was approximately 1.64 meters per second (m/s) or 3.66 miles per hour (mph); the maximum sustained wind speed was 8.31 m/s (18.6 mph) and the median wind speed was 1.43 m/s (3.2 mph).

The land within a one-mile radius of the LEHR facility is owned both privately and by UC Davis, and is used for animal research, agriculture and recreation. Immediately to the north, east and west of the LEHR facility are UC Davis research facilities. Privately-owned lands within one mile to the south and east of the LEHR Facility include permanent residences and some crop land. Approximately 75% of the surrounding land in the general vicinity of the LEHR facility is used for agriculture. Major crops include fruits, nuts and grains. Approximately 40% of the agricultural land in the vicinity is irrigated, and some of the nearby lands are used for cattle grazing (DOE, 1988).

The LEHR facility contains laboratory buildings and former animal-handling facilities (Figure 2). Approximately 45% of the Site is paved or covered by structures. Approximately 50% is unpaved and relatively free of vegetation, and 5% is covered by large, deep-rooted vegetation. Outdoor dog pens consisting of asphalt, gravel and soil occupy the remaining 5% of the LEHR facility. The land is owned by the Regents of the University of California and DOE owns most of the buildings on site.

In the early 1950s, the Atomic Energy Commission (now DOE) began conducting radiological studies at UC Davis on laboratory animals, particularly beagles. Initial studies were carried out on the main campus and involved the irradiation of beagles. DOE began operating in its present location in 1958 when full-scale experimental use of radioactive materials began. Research at the Site through the mid-1980s focused on the health effects from chronic exposure to radionuclides, primarily strontium-90 (Sr-90) and radium-226 (Ra-226). In the early 1970s, a cobalt-60 irradiator facility was constructed at the Site to study the effects of chronic exposure to gamma rays on bone marrow cells of beagles. In 1975, DOE initiated a program at the Site to study the potential health effects of combustion products from fossil fuel power plants. In 1983, the Toxic Pollutant Health Research Laboratory was established at the site. DOE-funded research at the LEHR facility ended in 1989. The LEHR facility is presently occupied by the UC Davis Center for Health and the Environment and the LEHR environmental restoration/waste management (ER/WM) project. UC Davis radionuclide emissions are not included in this report.

1.2 Source Description

The NESHAP requirements primarily target point source/stack emissions. However, a Memorandum of Understanding (MOU) between DOE and US EPA (DOE, 1995) applies the same point source criteria to potential non-point diffuse area sources.

1.2.1 Point Sources

No potential point sources of radionuclide emissions existed at the Site in 2002.

1.2.2 Non-Point Sources

Non-point diffuse radionuclide emission sources were limited to wind-blown fugitive dust from the following areas (Table 1):

- Western Dog Pens (WDPs) area;
- Eastern Dog Pens (EDPs) area;
- Domestic Septic System (DSS) 3; and
- DSS 6.

A removal action (RA) at the DSS 3 and 6 areas was conducted in 2002. Emissions from this RA are included in this report.

1.2.3 Western and Eastern Dog Pens Areas

The WDPs and EDPs areas, located near the center of the LEHR facility (Figure 2), are a potential non-point diffuse source of radiological emissions. The dog pens were used to house dogs involved in the Ra-226 and Sr-90 research activities at the Site. Excreta from dogs housed in outdoor pens contained low levels of radiological constituents. Solids were removed from the pens on a daily basis. Urine evaporated and/or infiltrated the gravel fill in the pens. Calculations by Marvin Goldman in a 1963 memorandum (DOE archived records) estimated that about 500 microCuries per year ($\mu\text{Ci}/\text{yr}$) of Sr-90 and about 50 $\mu\text{Ci}/\text{yr}$ of Ra-226 were eliminated in urine by the dogs housed in the pens (WA, 2000).

The WDPs area originally contained 320 pens. In 1975, 64 pens were removed to construct the Cellular Biology Lab (Building H-294, Figure 2). The EDPs area contained 96 pens and overlies UC Davis Landfill No. 2.

In 1996, the barrels that sheltered the dogs, interior chain-link fencing and concrete pedestals for the shelter were removed from the EDPs and existing 256 WDPs. The barrels and concrete pedestals were properly packaged and shipped to the DOE Hanford site for disposal. In 1999, the interior chain-link fencing was released according to DOE Order 5400.5 and recycled off site (EMS, 1999).

Between May and August 1 2001, the concrete curbing, gravel, metal grating and fence posts were removed from the existing 256 WDPs. Approximately 1,725 cubic yards (cu yd) of gravel,

650 cu yd of asphalt, 800 cu yd of concrete curbs, and 45 cu yd of metal grating and fence posts were removed. This material was removed using an excavator, segregated into one of four waste streams and transported to the appropriate stockpile on site using a wheel loader. Following the 2001 WDPs RA, the southern portion of the pens was backfilled to grade with clean imported fill.

Estimates of the WDPs area emissions are based on the maximum detected radionuclide concentrations that exceed background in soil samples collected from the upper two feet of soil in the WDPs. To be conservative, the data from all the WDPs Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) investigations (Table 2) were included in the data set. The emissions are assumed to be generated only from the northern portion of the WDPs (Figure 2) because the southern portion of the WDPs were covered with clean imported fill. The EDPs area emissions were based on the maximum detected radionuclide concentrations (Table 3) that exceed background in soil samples collected during the 1999 EDPs investigation, the only CERCLA investigation in this area.

1.2.4 Domestic Septic Systems 3 and 6

Beginning in 1958, the DSSs served LEHR offices and laboratories. A plan dated December 12, 1958 shows DSS 6, the oldest DSS at LEHR, connected to the "Animal Holding Building." DSS 3 was apparently installed during construction of the Reproductive Biology Laboratory (Building H-215) and the Specimen Storage Building (Building H-216) between 1962 and 1964, when the capacity of the tank serving the Inter-Regional Project No. 4 Building (Building H-217) may have become insufficient (WA, 2002a). The locations of DSSs 3 and 6 are shown on Figure 2.

Liquid wastes and sewage were discharged to DSSs 3 and 6 prior to the Site's connection to the UC Davis Wastewater Treatment Plant in 1971. At that time, Domestic Septic Tanks (DSTs) 3 and 6 were abandoned and no longer used. DSTs 3 and 6 were backfilled with sand and the influent/effluent lines for each tank were reportedly cut and capped (IT Corp., 1996). No formal closure reports for these DSTs are known to exist (Dames & Moore, 1994).

The RAs for DSSs 3 and 6 were conducted from April to July 2002. The DSS 3 RA began with demolition and removal of the distribution box (DB). All of the DB effluent lines were removed along with concrete, perforated Orangeburg pipe and leach trench gravel. Approximately one ft of additional soil from the trench floor and sidewalls was also removed. The DSS 3 leach trench excavation depth ranged from 11 to 12.5 ft below ground surface (bgs), and measured 8 ft wide and 50 ft long. Approximately 170 cu yd of piping, concrete, gravel and underlying soil were removed.

The DSS 6 RA consisted of removing of all the effluent lines, perforated Orangeburg pipe and leach trench gravel. Approximately one ft of additional soil from the trench floor and sidewalls was also removed. The excavation depth ranged from 6 ft to 7 ft bgs and was 11 ft wide by 105 ft long. Approximately 215 cu yd of piping, gravel and underlying soil were removed.

Ambient air samples were collected once during the RA for DSSs 3 and 6 from three stations around the perimeter of the LEHR facility and one distant background station (Air Monitoring Station [AM]-3, Figure 2). Details of the air monitoring results are presented in the *Draft DOE Areas Remedial Investigation Report* (WA, 2002b). Since there was only one sampling event, there were not adequate data to perform statistical tests. All of the maximum detected radionuclide concentrations in ambient air were below background Derived Concentration Guide (WA, 2002b). During the RA, particulate matter with an aerodynamic diameter less than or equal to 10 micrometers (PM₁₀) concentrations did not exceed the 150 micrograms per cubic meter (µg/m³) US EPA air quality standard for a 24-hour time period. The maximum PM₁₀ concentration detected on site during the RA was 20.6 µg/m³, collected from AM-7 located north of the WDPs area (Figure 2).

2. AIR EMISSION DATA

2.1 Point Sources

As discussed in Section 1, there are currently no point sources of radionuclide emissions at the Site.

2.2 Non-Point Sources

As discussed in Section 1, four potential non-point diffuse sources of radionuclide emissions were present at the Site in 2002. These included wind-blown, fugitive dust emission of radionuclides from the WDPs, EDPs, DSS 3 and DSS 6 areas. Based on the surface and shallow soil sampling results from the WDPs and EDPs, surface contamination is conservatively assumed to exist across the entire area of each potential radionuclide non-point emissions source. The 2002 DSSs 3 and 6 RA was a significant change to site conditions that could have resulted in emissions different from those estimated in the *Final Radionuclide Air Emission Annual Report for Calendar Year 2001* (WA, 2002c).

To calculate air emissions for this report, the surface area of each non-point source area was determined, using a scaled map of the Site, as follows:

- WDPs surface area = 5,263 square meters (m²) (the northern area of the WDPs that is not covered with clean fill; Figure 2);
- EDPs surface area = 3,900 m²;
- DSS 3 surface area = 60.6 m²; and,
- DSS 6 surface area = 115.2 m².

The particulate re-suspension rate model was used to calculate the fugitive dust emission rate based on the US EPA's guidance document *Rapid Assessment of Exposure to Particulate Emissions from Surface Contamination Sites* (Cowherd, 1985). Cowherd provides a methodology for the rapid, worst-case assessment of inhalation exposure to respirable particulate emissions, defined as airborne particles equal to or smaller than PM₁₀. PM₁₀ emission rates from site fugitive non-point sources were estimated using the equation developed by Cowherd for estimating respirable particle emissions from wind erosion of surfaces with an "unlimited reservoir" of erodible particles, adjusted for site-specific data using the following formula:

$$E_{10} = 0.036 \times (1 - V) \times \left(\frac{[u]}{u_t} \right)^3 \times F(x) \quad (\text{Eq. 1})$$

where

- E_{10} = annual average PM_{10} emission rate per unit contaminated surface, grams per square meter-hour [$g/(m^2 \cdot hr)$];
- V = fraction of contaminated surface vegetative cover (assumed zero for worst-case bare soil);
- $[u]$ = mean annual wind speed at 7 meters height = $[u]_{\text{site}} \times \ln(700/z_0) / \ln(300/z_0) = 1.94$ meters per second (m/s);
- $[u]_{\text{site}}$ = 95% upper confidence limit (UCL) of LEHR site meteorological tower mean annual wind speed at a height of three meters (= 1.66 m/s);
- z_0 = emission source area roughness height (= two centimeters [cm]);
- u_t = threshold value of wind speed at 7 meters high = $u_{\text{friction}} / 0.4 \times \ln(700/z_0) = 7.3$ m/s;
- u_{friction} = threshold friction velocity (= 50 cm per second [cm/s]);
- x = $0.866 u_t / [u]$ = dimensionless ratio; and,
- $F(x)$ = function plotted in Figure 4-3 (= 5.6×10^{-2}) of *Rapid Assessment of Exposure to Particulate Emissions from Surface Contamination Sites* (Cowherd, 1985).

The fraction of contaminated surface vegetative cover was conservatively assumed to be zero (i.e., bare soil) to calculate a worst-case value. The roughness height of two centimeters represents a value between the ranges for a plowed field and grasslands (Cowherd, 1985). Cowherd recommends a procedure for determining the threshold friction velocity based upon surface soil sieve analysis data. Because no surface soil sieve analysis data were available for the Site, a conservative threshold friction velocity of 50 cm/s was selected, following Cowherd's example (Application No. 1), for a rural emergency response application (Cowherd, 1985).

Cowherd recommends using the mean annual wind speed data tabulated in Table 4-1 of the particulate emission rate guidance document (Cowherd, 1985). However, since the tabulated values are based on 1977 meteorological data from Sacramento, and because site-specific data were available for 2002 from the LEHR on-site meteorological tower, the on-site data were selected as more representative of local site conditions. Therefore, the 95% UCL of the mean annual wind speed data was calculated using meteorological data collected during 2002 from the site meteorological tower.

The 95% UCL was calculated using equations for characterizing confidence limits of the mean for lognormal populations (Gilbert, 1987), as recommended by US EPA guidance (US EPA, 1992a). The resulting 95% UCL of the LEHR site meteorological tower mean annual wind speed for 2001 was 1.66 m/s. Because the height of the on-site meteorological tower is three meters, the

wind speed must be converted to the equivalent wind speed at a height of seven meters using a variation of Cowherd's Equation (see parameter [u] above), which assumes a logarithmic velocity profile near the Earth's surface.

Using Equation 1 and the data discussed above, the annual average PM₁₀ emission rate per unit contaminated surface was calculated as $E_{10} = 3.78 \times 10^{-5} \text{ g/m}^2\text{-hr}$. This emission rate is combined with the maximum radionuclide activity measured above background in surface soil activities for each potential emission source to calculate a particulate emission rate based on the following equation:

$$E_{\text{area}} = E_{10} \times A_{\text{area}} \times (24\text{-hr/day} \times 365 \text{ days/yr}) \times C_{\text{Rad}} \times (10^{-12} \text{ Curies per pCi})(\text{Eq. 2})$$

where,

- E_{area} = annual average PM₁₀ emission rate for the potential radionuclide emission non-point source, Curies per year (Ci/yr);
- E_{10} = annual average PM₁₀ emission rate per unit contaminated surface, g/(m²-hr);
- A_{area} = surface area of potential radionuclide emission non-point source, m²; and
- C_{Rad} = maximum value of the measured surface soil radionuclide activities, pCi/g.

Equations 1 and 2 and the methodology discussed above were used to estimate the non-point source fugitive dust radionuclide emission rate for each of the four potential radionuclide emission sources, as presented in Tables 2 through 5.

2.3 Particulate Matter-10

PM₁₀ data were calculated based on a volumetric flow rate. Glass fiber filters prepared by a US EPA-approved commercial laboratory were used to collect particulates greater than 10 μm at an average flow rate of 1.1 cubic meters (m³) per minute for approximately 24 hours. The total air volumes, V, ranged from 1,595 to 1,633 m³ based on the exposure time. PM₁₀ samples were collected from the four air monitoring station locations on May 22, 2002 (Figure 2). The concentration of PM₁₀ was calculated with the equation below:

$$C_{\text{PM-10}} = \frac{M_{\text{ff}} - M_{\text{fi}}}{V} \quad (\text{Eq. 3})$$

where,

- $C_{\text{PM-10}}$ = concentration of PM₁₀; μg/m³;
- M_{ff} = weight of the glass fiber filter after approximately 24 hours of exposure time reported by the laboratory; micrograms (μg);
- M_{fi} = weight of the glass fiber filter before sampling reported by the laboratory μg; and
- V = total air volume, m³.

3. DOSE ASSESSMENTS

3.1 Description of Dose Model

Compliance with the NESHAP requirements for diffuse, non-point source emissions was assessed using the US EPA atmospheric dispersion/radiation dose calculation computer code, CAP88-PC Version 1.0. This code was used to calculate the EDE to individual receptors at various distances from the four potential site radionuclide emission non-point sources. A total of four "individual receptor" CAP88-PC runs were executed to model the fugitive dust emission sources described in Section 2. For each of the four potential radionuclide emission non-point sources, a human receptor was identified in each of the north, south, east and west quadrants relative to the potential source.

The area source algorithm employed by CAP88-PC, Version 1.0, assumes the distance from an area source to a receptor is measured as the distance from the centroid of the area source to the receptor (US EPA, 1992b). For the site CAP88-PC modeling, the distance from an area source to a receptor is measured as the approximate distance from the centroid of the area source to the building assumed to house the receptor. Each CAP88-PC run included receptor distances to the respective source's maximally exposed individual (MEI), as well as the distances to MEIs identified for the other potential radionuclide emission sources ("Location" column for each source in Table 6).

The reported EDE to an MEI at the LEHR facility includes contributions from the four potential site radionuclide emission non-point sources based on the CAP88-PC model output. Based on the combined non-point source exposures, the MEI at the LEHR facility is located in the Inter-Regional Project No. 4 (Building H-217) (Figure 2, Table 6).

The collective population dose is calculated as the average radiation dose to an individual in a specified area, multiplied by the number of individuals in that area. Four "population" CAP88-PC runs were executed to model the four fugitive dust emission non-point sources. The CAP88-PC model output for each run is included in Appendix A. For each of the four potential radionuclide emission non-point sources, the CAP88-PC model was run with an updated population data file used in the *Radionuclide Air Emission Annual Report for Calendar Year 1998* (WA, 1999a), which includes the residential population in Davis and the student population at the UC Davis campus. The method used over the last five years to estimate population has been to take the 1997 data and add 2.4% per year (the average annual population growth for the City of Davis) (City of Davis web site: <http://www.city.davis.ca.us/aboutdavis/cityprofile/index.cfm?topic=5>).

The population file includes receptors within a distance of 10 kilometers (km) from the Site, rather than 80 km as specified in DOE guidance. This modification was necessary to avoid including the large number of receptors in the Sacramento area whose exposure to radionuclides resulting from the Site would be negligible, but whose population numbers would have a large effect on the collective population dose results. This approach is appropriate for calculating the collective population dose for the primarily rural LEHR facility surroundings. The results of the CAP88-PC population runs based upon the combined source exposures are presented in Table 7 and Appendix A.

Supplemental information required by DOE (DOE, 1994b) is included in Appendix B.

3.2 Summary of Input Parameters

The input parameters for the CAP88-PC runs are summarized for the WDPs, EDPs, and DSSs 3 and 6 areas in Tables 8, 9, 10 and 11, respectively. As noted above, the areal extent of each non-point source was conservatively calculated assuming that the maximum concentration (not corrected for background) of the observed radiological surface and shallow soil contamination for the WDPs, EDPs areas, and the maximum radionuclide concentrations from investigation and waste characterization sampling for the DSSs 3 and 6 areas, were present across the entire potential radionuclide emission non-point source area. Conservative radionuclide emissions were calculated using the maximum of the soil radionuclide concentrations measured for each potential area non-point source and applying the US EPA-recommended particulate re-suspension rate model to calculate the fugitive dust emission rates.

The Sacramento area wind file included with the CAP88-PC computer code was used for the modeling. Site meteorological data were not utilized for the CAP88-PC modeling due to incompatible formats between on-site data (MET144) and the CAP88-PC requirement (CD144). Use of the Sacramento wind file is appropriate because of the Site's proximity (approximately 15 miles) to Sacramento, the similar geography of the two areas, the lack of intervening geographical anomalies, and the absence of a compatible meteorological data file from a closer air station.

4. COMPLIANCE ASSESSMENT

Point Source Effective Dose Equivalent: None

Non-Point Source Maximum Effective Dose Equivalent: 3.8×10^{-4} millirem per year [mrem/yr]
 $(3.8 \times 10^{-6}$ milliSievert [mSv]/yr)

Location of On-Site Maximally Exposed Individual: Inter-Regional Project No. 4 (H-217), 34m north of DSS 3

4.1 Certification

I certify under penalty of law that I have personally examined and am familiar with the information submitted herein and based on my inquiry of those individuals immediately responsible for obtaining the information, I believe that the submitted information is true, accurate and complete. I am aware that there are significant penalties for submitting false information including the possibility of fine and imprisonment (See 18 U. S. C. 1001).

Signature: Robert D. Devany Date: 4-16-03
Robert Devany
LEHR Site ER/WM Project Manager

Signature: Jay B. Tomlin Date: 4-16-03
Jay Tomlin
DOE-LEHR Project Manager

5. ADDITIONAL INFORMATION

In 2002, the LEHR facility completed construction or modifications in the DSSs 3 and 6 areas. The facility is exempt from filing an application to construct or modify per the MOU (DOE, 1995) because the EDE caused by all emissions from the completed modifications and construction is less than 1% of the standard in 40 CFR 61.92, 10 mrem/yr.

As required in the MOU for facilities that were exempt from having to submit an application to US EPA for construction or modifications, the following information is provided for the Site:

- **Provide a brief description of the construction or modification project and an estimate of potential doses to the public.**

There were construction or modification projects completed in 2002 at the LEHR facility for which approval to construct or modify was required or waived under Section 61.96 of Subpart H of 40 CFR 61. Between April and July of 2002, contaminated concrete, Orangeburg pipe and leach trench gravel were excavated and removed from the DSS 3 and 6 areas for proper storage and disposal. Approximately 170 and 215 cu yd of material were removed from the DSS 3 and 6 areas, respectively.

The collective population equivalent dose from the DSS 3 and 6 RAs to Davis residents was $4.44E \times 10^{-5}$ person-roentgen equivalent man per year, and the EDE for the off-site MEI was 8.66×10^{-6} mrem/yr, as estimated by CAP88-PC. The predicted EDE for the off-site MEI is several orders of magnitude below the 10 mrem/yr standard required by 40 CFR Part 61 Subpart H.

- **Identify any unplanned releases of radionuclides to the atmosphere.**

There were no unplanned releases of radionuclides to the atmosphere during 2002.

- **Results of the dose assessment associated with the diffuse source emissions from the facility.**

As noted in Sections 1 and 2, there are currently no point sources of radionuclide emissions at the Site. The only potential sources of radionuclide emissions remaining at the Site are diffuse sources of fugitive dust, and the results of the dose assessment associated with the diffuse non-point source emissions from the Site are presented in Section 3 and Table 4. The total contribution to the EDE to the on-site MEI resulting from non-point source

emissions was estimated to be 3.8×10^{-4} mrem/yr (3.9×10^{-6} mSv/yr), far below the 10 mrem/yr standard.

6. REFERENCES

- City of Davis Website, <http://www.city.davis.ca.us/aboutdavis/cityprofile/index.cfm?topic=5>.
- Cowherd, C., 1985, *Rapid Assessment of Exposure to Particulate Emissions from Surface Contamination Sites*, U.S. Environmental Protection Agency, Washington, D.C. (USEPA/600/8-85/002).
- United States Department of Energy (DOE), 1988, *Environmental Survey Preliminary Report*, Laboratory for Energy-Related Health Research, Davis, California, Environment, Safety and Health Office of Environmental Audit.
- Dames & Moore (D&M), 1994, Remedial Investigation, Feasibility Study and Environmental Assessment Work Plan, LEHR Environmental Restoration, University of California, Davis.
- DOE, 1990, *Results of the U.S. Department of Energy Indoor Radon Study*, August 1990.
- DOE, 1994a, *Water Monitoring Plan*, LEHR Environmental Restoration, UC Davis, California.
- DOE, 1994b, *Memorandum—Calendar Year 1993 Radionuclide Air Emissions Annual Reports for DOE Sites*, March.
- DOE, 1995, *Memorandum of Understanding with the Environmental Protection Agency Concerning the Radionuclide National Emission Standards for Hazardous Air Pollutants*, U.S. Department of Energy Memorandum; April 5, 1995.
- DOE, Various DOE Records Archived at the Federal Records Center, 1000 Commodore Drive, San Bruno, California.
- Environmental Management Services (EMS), 1999, *Final Miscellaneous Metal Disposition Report for the Laboratory for Energy-Related Health Research*, University of California, Davis, Rev. 0, May.
- Gilbert, Richard O., 1987, *Statistical Methods for Environmental Pollution Monitoring; Section 13.2: Characterizing Lognormal Populations: Confidence Limits of the Mean*, Van Nostrand Reinhold, New York, 1987.
- IT Corporation (IT Corp.), 1996, *Limited Field Investigation Work Plan, LEHR*, University of California, Davis.
- United States Environmental Protection Agency (US EPA), 1992a, *Supplemental Guidance to RAGS: Calculating the Concentration Term*, U.S. Environmental Protection Agency (PB92-963373, May 1992).
- US EPA, 1992b, *User's Guide for CAP88-PC Version 1*, U.S. Environmental Protection Agency, (402-3-92-001, March 1992).

- WA, 1999a, *Final Radionuclide Air Emission Annual Report (Subpart H of 40 CFR 61) Calendar Year 1998 for the Laboratory for Energy-Related Health Research*, University of California, Davis, June, Rev. 0.
- WA, 2000, *Draft Final Engineering Evaluation/Cost Analysis for the Western and Eastern Dog Pens at the Laboratory for Energy-Related Health Research*, University of California, Davis, November, Rev. E.
- WA, 2002a, *Domestic Septic Systems 3 and 6 Removal Actions Work Plan for Laboratory for Energy-Related Health Research*, University of California, Davis, May, Rev. 0.
- WA, 2002b, *Draft DOE Areas Remedial Investigation Report for the Laboratory for Energy-Related Health Research*, University of California, Davis, March, Rev. C.
- WA, 2002c, *Final Radionuclide Air Emission Annual Report (Subpart H of 40 CFR 61) Calendar Year 2000 for the Laboratory for Energy-Related Health Research*, University of California, Davis, May, Rev. 0.

TABLES

Table 1. Summary of LEHR Site Potential Non-Point Diffuse Area Radionuclide Sources

Potential Fugitive Source	Description
Western Dog Pens Area	Formerly housed animals involved in LEHR facility research. Although the area was remediated in 2001, a portion of WDPs soil remains uncovered.
Eastern Dog Pens Area	Formerly housed animals involved in LEHR facility research.
Domestic Septic Systems 3 Area	Subsurface tanks and leachfield which received sink and lavatory waste from research facilities. The leachfield was remediated in 2002.
Domestic Septic System 6 Area	Subsurface tanks and leachfield which received sink and lavatory waste from research facilities. The leachfield was remediated in 2002.

Abbreviation

LEHR Laboratory for Energy-Related Health Research
DSS Domestic Septic System

Table 2. Summary of Maximum Surface Soil Concentrations for Radionuclides Exceeding Background and PM₁₀ Emission Rates, LEHR Western Dog Pens Area

Radionuclide	Maximum Concentration ¹ (pCi/g)	Maximum PM ₁₀ Emission Rate (Ci/yr)
Actinium-228	0.719	1.25E-09
Bismuth-212	0.628	1.1E-09
Bismuth-214	1.09	1.90E-09
Carbon-14	11.3	1.97E-08
Cesium-137	0.115	2.01E-10
Cobalt-60	0.028	4.88E-11
Lead-210	4.96	8.65E-09
Lead-212	0.744	1.30E-09
Lead-214	1.41	2.46E-09
Potassium-40	14.3	2.49E-08
Radium-226	1.9	3.31E-09
Strontium-90	0.675	1.18E-09
Thallium-208	0.272	4.74E-10
Thorium-234	1.62	2.83E-09
Uranium-235	0.232	4.05E-10
Uranium-238	1.62	2.83E-09

Notes

¹Not corrected for background.

Abbreviations

Ci/yr Curies per year

pCi/g picoCuries per gram

PM₁₀ Particulate matter with an aerodynamic diameter less than or equal to 10 micrometers

Table 3. Summary of Maximum Surface Soil Concentrations for Radionuclides Exceeding Background and PM₁₀ Emission Rates, LEHR Eastern Dog Pens Area

Radionuclide	Maximum Concentration ¹ (pCi/g)	Maximum PM ₁₀ Emission Rate (Ci/yr)
Bismuth-212	0.415	5.36E-10
Bismuth-214	0.572	7.39E-10
Cesium-137	0.191	2.47E-10
Lead-214	0.607	7.85E-10
Strontium-90	0.164	2.12E-10
Thallium-208	0.219	2.83E-10
Thorium-228	1.54	1.99E-09
Thorium-232	1.26	1.63E-09
Thorium-234	0.89	1.15E-09
Tritium	1.21	1.56E-09
Uranium-235	0.383	4.95E-11

Notes

¹Not corrected for background.

Abbreviations

Ci/yr Curies per year

pCi/g picoCuries per gram

PM₁₀ Particulate matter with an aerodynamic diameter less than or equal to 10 micrometers

Table 4. Summary of Maximum Surface Soil Concentrations for Radionuclides Exceeding Background and PM₁₀ Emission Rates, LEHR Domestic Septic System 3

Radionuclide	Maximum Concentration ¹ (pCi/g)	Maximum PM ₁₀ Emission Rate (Ci/yr)
Bismuth-214	2.18	4.38E-11
Carbon-14	0.155	3.11E-12
Cesium-137	0.0619	1.24E-12
Lead-210	2.15	4.32E-11
Lead-214	2.33	4.68E-11
Plutonium-241	0.664	1.33E-11
Radium-226	2.44	4.90E-11
Strontium-90	2.01	4.04E-11
Thorium-234	0.859	1.73E-11
Uranium-233/234	1.1	2.21E-11
Uranium-235/236	0.0903	1.81E-12
Uranium-238	0.649	1.30E-11

Notes

¹Not corrected for background.

Abbreviations

Ci/yr Curies per year

pCi/g picoCuries per gram

PM₁₀ Particulate matter with an aerodynamic diameter less than or equal to 10 micrometers

Table 5. Summary of Maximum Surface Soil Concentrations for Radionuclides Exceeding Background and PM₁₀ Emission Rates, LEHR Domestic Septic System 6

Radionuclide	Maximum Concentration (pCi/g)	Maximum PM ₁₀ Emission Rate (Ci/yr)
Actinium-228	0.67	2.56E-11
Americium-241	0.0184	7.03E-13
Bismuth-212	0.45	1.72E-11
Bismuth-214	0.61	2.33E-11
Cesium-137	0.0549	2.10E-12
Cobalt-60	0.013	4.96E-13
Lead-210	1.75	6.68E-11
Lead-214	0.75	2.86E-11
Strontium-90	0.211	8.06E-12
Thallium-208	0.242	9.24E-12
Tritium	2.3	8.78E-11
Uranium-233/234	0.737	2.81E-11
Uranium-235/236	0.0449	1.71E-12

Abbreviations

Ci/yr Curies per year

pCi/g picoCuries per gram

PM₁₀ Particulate matter with an aerodynamic diameter less than or equal to 10 micrometers

Table 6. Summary of On-Site Effective Dose Equivalent to Maximally Exposed Individual Resulting from Radionuclide Emissions from Each Potential Fugitive Dust Emission Non-Point Source

MEI Receptor Description	<u>Western Dog Pens Area</u>		<u>Eastern Dog Pens Area</u>		<u>Domestic Septic System 3</u>		<u>Domestic Septic System 6</u>		Maximum Total Dose (mrem/yr) ³
	(mrem/yr) ¹	Location ²	(mrem/yr) ¹	Location ²	(mrem/yr) ¹	Location ²	(mrem/yr) ¹	Location ²	
Specimen Storage Building (Building H-216)	2.7E-04	48 m W	5.6E-05	132 m W	3.20E-05	15 m N	1.30E-06	58 m S	3.6E-04
UC Davis Building E of LEHR Site	1.9E-05	300 m E	8.7E-06	180 m E	3.80E-07	346 m E	2.00E-07	338 m E	2.8E-05
Off-Site Receptor S of Putah Creek	1.7E-05	1,200 m S	6.3E-07	1,000 m S	3.70E-07	1173 m S	1.80E-07	1246 m S	1.8E-05
Off-Site Receptor W of LEHR Site	2.2E-05	400 m W	4.2E-06	500 m W	4.00E-07	354 m W	2.30E-07	360 m W	2.7E-05
Animal Hospital Building No. 1 (Building H-219)	2.1E-04	65 m W	3.6E-05	165 m W	1.60E-06	85 m NW	8.50E-06	31m NW	2.6E-04
Inter-Regional Project No. 4 Building (Building H-217)	3.2E-04	52 m W	4.8E-05	143 m W	7.30E-06	34 m N	1.80E-06	43 m S	3.8E-04
Animal Hospital Building No. 2 (Building H-218)	2.1E-04	65 m W	3.6E-05	165 m W	3.30E-06	53 m NW	2.70E-06	31 m NW	2.5E-04
Cellular Biology Laboratory (Building H-294)	2.7E-04	65 m N	4.2E-05	150 m NNE	8.20E-07	118 m NE	2.60E-06	52 m NE	3.2E-04
Clinical Pathology (H-215)	1.00E-04	99 m W	4.40E-05	150 m W	1.40E-05	24 m N	7.60E-07	61 m SW	1.6E-04
Main Office (H-213)	2.70E-04	65 m NW	3.80E-05	187 m NW	1.20E-06	110 m N	4.50E-06	43 m N	3.1E-04

Notes

¹ The EDE to the MEI is taken as the maximum modeled dose within a 45° sector in the direction and at the distance indicated in the “location” column. For example, the dose 65 m north of the Western Dog Pens Area would be the maximum modeled dose at 65 m N, 65 m NNE and 65 m NNW.

² The distance from an area source to a receptor is defined by CAP88-PC as the distance from the centroid of the area source to the receptor (US EPA, 1992b). For the LEHR facility CAP88-PC modeling, the distance from an area non-point source to a receptor is measured as the approximate distance from the centroid of the area non-point source to the centroid of the building assumed to house the receptor.

Table 6. Summary of On-Site Effective Dose Equivalent to Maximally Exposed Individual Resulting from Radionuclide Emissions from Each Potential Fugitive Dust Emission Non-Point Source (continued)

³ The maximum total dose is the sum of EDEs modeled for each MEI receptor from the four potential radionuclide fugitive dust emission non-point sources. Value in boldface is the maximum total dose for the site-wide MEI.

Abbreviations

EDE	effective dose equivalent
E	east
m	meters
MEI	Maximally Exposed Individual
mrem/yr	millirem per year
N	north
NNE	north by northeast
NNW	north by northwest
No.	number
S	south
UC Davis	University of California, Davis
W	west

Table 7. Summary of Estimated Collective Population Dose Resulting from Radionuclide Emissions from Each Fugitive Dust Emission Non-Point Source

Potential Emission Source	Off-Site Maximally Exposed Individual		Collective Population Dose (person-rem/yr)
	(mrem/yr)	Location	
Western Dog Pens Area	4.53E-06	250 m north ¹	1.43E-05
Eastern Dog Pens Area	2.7E-05	250 m north ¹	8.2E-05
Domestic Septic System 3	1.76E-07	250 m north ¹	5.47E-07
Domestic Septic System 6	1.70E-07	250 m north ¹	5.30E-07
Total LEHR Site	3.16E-05		9.7E-05

Notes

Source of data: CAP88-PC Version 1.0 modeling output files.

The collective population dose is for receptors within a 10-km radius.

¹The approximate location of UC Davis research facilities.

Abbreviations

km kilometer(s)
 m meter(s)
 mrem/yr millirem per year
 N north
 person-rem/yr person-roentgen equivalent man per year
 UC Davis University of California, Davis

Table 8. CAP88-PC Inputs for the Western Dog Pens Area

Run type	Individual	Receptor distances (m): 48, 52, 65, 99, 300, 400, 1200			
Run type	Population	Population file to use: 02LEHR.POP			
Modeling Options	Generate genetic effects?	Yes			
	Create Dose and Risk Factor file?	Yes			
	Create Concentration Table file?	Yes			
	Create Chi/Q Table file?	Yes			
Meteorological Data					
Wind file to use:	SAC0320.WND (CAP88-PC supplied wind file)				
Annual precipitation (cm/yr):	26.39 (2002 Davis total)				
Annual ambient temperature (°C):	14.9 (2002 Davis average)				
Height of lid (m):	1,000 (CAP88-PC default)				
Source Data					
Source type:	AREA				
Number of source:	1				
Height (m):	0				
Area (m ²):	5,263				
Plume rise:	Zero (Plume rise is zero for each Pasquill stability category)				
Agricultural Data					
	Source: Rural (CAP88-PC defaults)				
	Vegetable	Milk	Beef		
Fraction home produced:	0.7	0.399	0.442		
Fraction from assessment area:	0.3	0.601	0.558		
Fraction imported:	0	0	0		
Beef cattle density (#/km ²):	8.81E-02 (CAP88-PC default)				
Milk cattle density (#/km ²):	2.85E-02 (CAP88-PC default)				
Land fraction cultivated for vegetable crops:	0.25 (Site specific parameter per 1995 NESHAP)				
Radionuclide Release Data					
PM ₁₀ Emission rate (g/m ² -sec):	3.78E-5 (Using Cowherd, 1985, unlimited erosion potential)				
	Nuclide ID	Max. Soil (pCi/g)	Emis. Rate (Ci/yr)	Size ¹ (µm)	Class ¹
	Ac-228	0.719	1.25E-09	1	Y
	Bi-212	0.628	1.10E-09	1	W
	Bi-214	1.09	1.90E-09	1	W
	C-14	11.3	1.97E-08	0	G
	Cs-137	0.115	2.01E-10	1	D
	Co-60	0.028	4.88E-11	1	Y
	Pb-210	4.96	8.65E-09	1	D
	Pb-212	0.744	1.30E-09	1	D
	Pb-214	1.41	2.46E-09	1	D
	K-40	14.3	2.49E-08	1	D
	Ra-226	1.9	3.31E-09	1	W
	Sr-90	0.675	1.18E-09	1	D
	Tl-208	0.272	4.74E-10	1	D
	Th-234	1.62	2.83E-09	1	Y
	U-235	0.232	4.05E-10	1	Y
	U-238	1.62	2.83E-09	1	Y

Table 8. CAP88-PC Inputs for the Western Dog Pens Area (continued)

Note

¹ CAP88-PC default particle size and lung retention class.

Abbreviations

km ²	per square kilometer
°C	degrees Celsius
Ac-228	actinium-228
Bi-212	bismuth-212
Bi-214	bismuth-214
C-14	carbon-14
Ci/yr	Curies per year
cm/yr	centimeter(s) per year
Co-60	cobalt-60
Cs-137	cesium-137
Emis.	emission
g/m ² -sec	grams per square meter-seconds
ID	identification (number)
K-40	potassium-40
m	meters
m ²	square meter(s)
Max.	maximum (concentration)
NESHAP	National Emissions Standards for Hazardous Air Pollutants
Pb-210	lead-210
Pb-212	lead-212
Pb-214	lead-214
pCi/g	picoCurie per gram
PM ₁₀	particulate matter with an aerodynamic diameter less than or equal to 10 micrometers
Ra-226	radium-226
Sr-90	strontium-90
Th-234	thorium-234
Tl-208	thallium-208
U-235	uranium-235
U-238	uranium-238
µm	micrometer

Table 9. CAP88-PC Inputs for the Eastern Dog Pens Area

Run type	Individual	Receptor distances (m): 132, 143, 150, 165, 180, 187, 500, 1000				
Run type	Population	Population file to use: 02LEHR.POP				
Modeling Options	Generate genetic effects?	Yes				
	Create Dose and Risk Factor file?	Yes				
	Create Concentration Table file?	Yes				
	Create Chi/Q Table file?	Yes				
Meteorological Data						
	Wind file to use:	SAC0320.WND (CAP88-PC supplied wind file)				
	Annual precipitation (cm/yr):	26.39 (2002 Davis total)				
	Annual ambient temperature (°C):	14.9 (2002 Davis average)				
	Height of lid (m):	1,000 (CAP88-PC default)				
Source Data						
	Source type:	AREA				
	Number of source:	1				
	Height (m):	0				
	Area (m ²):	3,900				
	Plume rise:	Zero (Plume rise is zero for each Pasquill stability category)				
Agricultural Data						
	Source:	Rural (CAP88-PC defaults)				
		Vegetable	Milk	Beef		
	Fraction home produced:	0.7	0.399	0.442		
	Fraction from assessment area:	0.3	0.601	0.558		
	Fraction imported:	0	0	0		
	Beef cattle density (#/km ²):	8.81E-02 (CAP88-PC default)				
	Milk cattle density (#/km ²):	2.85E-02 (CAP88-PC default)				
	Land fraction cultivated for vegetable crops:	0.25 (Site specific parameter per 1995 NESHAP)				
Radionuclide Release Data						
	PM ₁₀ Emission Rate (g/m ² -hr):	3.78E-5 (Using Cowherd, 1985, unlimited erosion potential)				
		Nuclide ID	Max. Soil (pCi/g)	Emis. Rate (Ci/yr)	Size ¹ (µm)	Class ¹
		Bi-212	0.415	5.36E-10	1	W
		Bi-214	0.572	7.39E-10	1	W
		Cs-137	0.191	2.47E-10	1	D
		Pb-214	0.607	7.85E-10	1	D
		Sr-90	0.164	2.12E-10	1	D
		Tl-208	0.219	2.83E-10	1	D
		Th-228	1.54	1.99E-09	1	Y
		Th-232	1.26	1.63E-09	1	Y
		Th-234	0.89	1.15E-09	1	Y
		H-3	1.21	1.56E-09	0	G
		U-235	0.0383	4.95E-11	1	Y

Table 9. CAP88-PC Inputs for the Eastern Dog Pens Area (continued)

Note

¹ CAP88-PC default particle size and lung retention class.

Abbreviations

km ²	per square kilometer
°C	degrees Celsius
Bi-212	bismuth-212
Bi-214	bismuth-214
Ci/yr	Curies per year
cm/yr	centimeter(s) per year
Cs-137	cesium-137
Emis.	emission
g/m ² -sec	grams per square meter-seconds
H-3	tritium
m	meter(s)
m ²	square meter(s)
Max.	maximum (concentration)
NESHAP	National Emissions Standards for Hazardous Air Pollutants
Pb-214	lead-214
pCi/g	picoCurie per gram
PM ₁₀	particulate matter with an aerodynamic diameter less than or equal to 10 micrometers
Sr-90	strontium-90
Th-228	thorium-228
Th-234	thorium-234
Tl-208	thallium-208
U-235	uranium-235
µm	micrometer

Table 10. CAP88-PC Inputs for the Domestic Septic System 3

Run type	Individual	Receptor distances (m): 15, 24, 34, 53, 85, 110, 118, 346, 1173, 354				
Run type	Population	Population file to use: 02LEHR.POP				
Modeling Options	Generate genetic effects?	Yes				
	Create Dose and Risk Factor file?	Yes				
	Create Concentration Table file?	Yes				
	Create Chi/Q Table file?	Yes				
Meteorological Data						
	Windfile to use:	SAC0320.WND (CAP88-PC supplied wind file)				
	Annual precipitation (cm/yr):	26.39 (2002 Davis total)				
	Annual ambient temperature (°C):	14.9 (2002 Davis average)				
	Height of lid (m):	1,000 (CAP88-PC default)				
Source Data						
	Source type:	AREA				
	Number of source:	1				
	Height (m):	0				
	Area (m ²):	652				
	Plume rise:	Zero (Plume rise is zero for each Pasquill stability category)				
Agricultural Data						
	Source:	Rural (CAP88-PC defaults)				
		Vegetable	Milk	Beef		
	Fraction home produced:	0.7	0.399	0.442		
	Fraction from assessment area:	0.3	0.601	0.558		
	Fraction imported:	0	0	0		
	Beef cattle density (#/km ²):	8.81E-02 (CAP88-PC default)				
	Milk cattle density (#/km ²):	2.85E-02 (CAP88-PC default)				
	Land fraction cultivated for vegetable crops:	0.25 (Site specific parameter per 1995 NESHAP)				
Radionuclide Release Data						
	PM ₁₀ Emission Rate (g/m ² -hr):	3.78E-5 (Using Cowherd, 1985, unlimited erosion potential)				
		Nuclide ID	Max. Soil (pCi/g)	Emis. Rate (Ci/yr)	Size ¹ (µm)	Class ¹
		Bi-214	2.18	4.38E-11	1	W
		C-14	0.155	3.11E-12	1	G
		Cs-137	0.0619	1.24E-12	1	D
		Pb-210	2.15	4.32E-11	1	D
		Pb-214	2.33	4.68E-11	1	D
		Pu-241	0.664	1.33E-11	1	D
		Ra-226	2.44	4.90E-11	1	W
		Sr-90	2.01	4.04E-11	1	D
		Th-233	0.859	1.73E-11	1	Y
		U-233 ²	1.1	2.21E-11	1	Y
		U-235 ³	0.0903	1.81E-12	1	Y
		U-238	0.649	1.30E-11	1	Y

Table 10. CAP88-PC Inputs for the Domestic Septic System 3 (continued)

Note

¹ CAP88-PC default particle size and lung retention class.

² Reported by analytical lab as U-233/234.

³ Reported by analytical lab as U-235/236.

Abbreviations

#/km ²	number per square kilometer
°C	degrees Celsius
Bi-214	bismuth-214
C-14	carbon 14
Ci/yr	Curies per year
cm/yr	centimeter(s) per year
Cs-137	cesium-137
Emis.	emission
g/m ² -sec	grams per square meter-seconds
m	meter(s)
m ²	square meter(s)
Max.	maximum (concentration)
NESHAP	National Emissions Standards for Hazardous Air Pollutants
Pb-214	lead-214
pCi/g	picoCurie per gram
PM ₁₀	particulate matter with an aerodynamic diameter less than or equal to 10 micrometers
Pu-241	plutonium 241
Ra-226	radium 226
Sr-90	strontium-90
Th-234	thorium-234
U-233	uranium-233
U-235	uranium-235
µm	micrometer

Table 11. CAP88-PC Inputs for the Domestic Septic System 6

Run type	Individual	Receptor distances (m): 43, 52, 58, 61, 338, 360, 1246			
Run type	Population	Population file to use: 02LEHR.POP			
Modeling Options	Generate genetic effects?	Yes			
	Create Dose and Risk Factor file?	Yes			
	Create Concentration Table file?	Yes			
	Create Chi/Q Table file?	Yes			
Meteorological Data					
Wind file to use:	SAC0320.WND (CAP88-PC supplied wind file)				
Annual precipitation (cm/yr):	26.39 (2002 Davis total)				
Annual ambient temperature (°C):	14.9 (2002 Davis average)				
Height of lid (m):	1,000 (CAP88-PC default)				
Source Data					
Source type:	AREA				
Number of source:	1				
Height (m):	0				
Area (m ²):	1,240				
Plume rise:	Zero (Plume rise is zero for each Pasquill stability category)				
Agricultural Data					
	Source: Rural (CAP88-PC defaults)				
	Vegetable	Milk	Beef		
Fraction home produced:	0.7	0.399	0.442		
Fraction from assessment area:	0.3	0.601	0.558		
Fraction imported:	0	0	0		
Beef cattle density (#/km ²):	8.81E-02 (CAP88-PC default)				
Milk cattle density (#/km ²):	2.85E-02 (CAP88-PC default)				
Land fraction cultivated for vegetable crops:	0.25 (Site specific parameter per 1995 NESHAP)				
Radionuclide Release Data					
PM ₁₀ Emission Rate (g/m ² -hr):	3.78E-5 (Using Cowherd, 1985, unlimited erosion potential)				
	Nuclide ID	Max. Soil (pCi/g)	Emis. Rate (Ci/yr)	Size ¹ (µm)	Class ¹
	Ac-228	0.67	2.56E-11	1	Y
	Am-241	0.0184	7.03E-13	1	W
	Bi-212	0.45	1.72E-11	1	W
	Bi-214	0.61	2.33E-11	1	W
	Cs-137	0.0549	2.10E-12	1	D
	Co-60	0.013	4.96E-13	1	Y
	Pb-210	1.75	6.68E-11	1	D
	Pb-214	0.75	2.86E-11	1	D
	Sr-90	0.211	8.06E-12	1	D
	Tl-208	0.242	9.24E-12	1	D
	H-3	2.3	8.78E-11	0	G
	U-233	0.737	2.81E-11	1	Y
	U-235	0.0449	1.71E-12	1	Y

Table 11. CAP88-PC Inputs for the Domestic Septic System 6 (continued)

Note

¹ CAP88-PC default particle size and lung retention class.

² Reported by analytical lab as U-233/234.

³ Reported by analytical lab as U-235/236.

Abbreviations

km ²	square kilometer
°C	degrees Celsius
Ac-228	actinium-228
Am-241	americium-241
Bi-214	bismuth-214
C-14	carbon-14
Ci/yr	Curies per year
cm/yr	centimeter(s) per year
Cs-137	cesium-137
Co-60	cobalt-60
Emis.	emission
g/m ² -sec	grams per square meter-seconds
m	meter(s)
m ²	square meter(s)
Max.	maximum (concentration)
NESHAP	National Emissions Standards for Hazardous Air Pollutants
Pb-210	lead-210
Pb-214	lead-214
pCi/g	picoCurie per gram
PM ₁₀	particulate matter with an aerodynamic diameter less than or equal to 10 micrometers
PU-241	plutonium-241
Ra-226	radium 226
Sr-90	strontium-90
Tl-208	thallium-208
U-233	uranium-233
U-235	uranium-235
µm	micrometer

FIGURES




Figure 1. LEHR Facility Location Map, UC Davis, California

Weiss Associates



Figure 2. LEHR Facility Potential and Known DOE Non-Point Source Areas

APPENDIX A

CAP88-PC OUTPUT RESULTS

H.syn

C A P 8 8 - P C

Version 1.00

Clean Air Act Assessment Package - 1988

S Y N O P S I S R E P O R T

Non-Radon Individual Assessment
Mar 5, 2003 1:39 pm

Facility: LEHR
Address: Old Davis Road
City: Davis
State: CA Zip: 95616

Effective Dose Equivalent
(mrem/year)

4.76E-04

At This Location: 48 Meters North Northwest
Source Category: nonpoint
Source Type: Area
Emission Year: 2002

Comments:

Dataset Name: WDPind
Dataset Date: Mar 5, 2003 1:35 pm
Wind File: WNDFILES\SAC0320.WND

Mar 5, 2003 1:39 pm

SYNOPSIS
Page 1

MAXIMALLY EXPOSED INDIVIDUAL

H.syn

Location Of The Individual: 48 Meters North Northwest
Lifetime Fatal Cancer Risk: 5.64E-09

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Dose Equivalent (mrem/y)
GONADS	7.55E-05
BREAST	7.26E-05
R MAR	3.03E-04
LUNGS	1.72E-03
THYROID	7.34E-05
ENDOST	3.50E-03
RMNDR	3.21E-04
EFFEC	4.76E-04

□

Mar 5, 2003 1:39 pm

SYNOPSIS
Page 2

RADIONUCLIDE EMISSIONS DURING THE YEAR 2002

Nuclide	Class	Size	Source #1 Ci/y	TOTAL Ci/y
AC-228	Y	1.00	1.2E-09	1.2E-09
BI-212	W	1.00	1.1E-09	1.1E-09
BI-214	W	1.00	1.9E-09	1.9E-09
C-14	*	0.00	2.0E-08	2.0E-08
CO-60	Y	1.00	2.0E-10	2.0E-10
CS-137	D	1.00	4.9E-11	4.9E-11
PB-210	D	1.00	8.6E-09	8.6E-09
PB-212	D	1.00	1.3E-09	1.3E-09
PB-214	D	1.00	2.5E-09	2.5E-09
K-40	D	1.00	2.5E-08	2.5E-08
RA-226	W	1.00	3.3E-09	3.3E-09
SR-90	D	1.00	1.2E-09	1.2E-09
TL-208	D	1.00	4.7E-10	4.7E-10
TH-234	Y	1.00	2.8E-09	2.8E-09
U-235	Y	1.00	4.1E-10	4.1E-10
U-238	Y	1.00	2.8E-09	2.8E-09

SITE INFORMATION

Temperature: 15 degrees C
Precipitation: 26 cm/y
Mixing Height: 1000 m

SOURCE INFORMATION

Source Number: 1

Source Height (m): 0.00
 Area (sq m): 5.26E+03

Plume Rise
 Pasquill Cat:

	A	B	C	D	E	F	G
Zero:	0.00	0.00	0.00	0.00	0.00	0.00	0.00

AGRICULTURAL DATA

	Vegetable	Milk	Meat
Fraction Home Produced:	0.700	0.399	0.442
Fraction From Assessment Area:	0.300	0.601	0.558
Fraction Imported:	0.000	0.000	0.000

Food Arrays were not generated for this run.
 Default Values used.

DISTANCES USED FOR MAXIMUM INDIVIDUAL ASSESSMENT

48 300 1200 400 65 52 99

I.syn

C A P 8 8 - P C

Version 1.00

Clean Air Act Assessment Package - 1988

S Y N O P S I S R E P O R T

Non-Radon Population Assessment
Mar 5, 2003 1:39 pm

Facility: LEHR
Address: Old Davis Road
City: Davis
State: CA Zip: 95616

Effective Dose Equivalent
(mrem/year)

2.07E-05

At This Location: 250 Meters North
Source Category: nonpoint
Source Type: Area
Emission Year: 2002

Comments:

Dataset Name: WDPpop
Dataset Date: Mar 5, 2003 1:39 pm
Wind File: WNDFILES\SAC0320.WND
Population File: POPFILES\02LEHR.POP

Mar 5, 2003 1:39 pm

SYNOPSIS
Page 1

MAXIMALLY EXPOSED INDIVIDUAL

Page 1

I.syn

Location Of The Individual: 250 Meters North
Lifetime Fatal Cancer Risk: 2.43E-10

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Selected Individual (mrem/y)	Collective Population (person-rem/y)
GONADS	3.00E-06	9.50E-06
BREAST	2.87E-06	9.09E-06
R MAR	1.26E-05	4.03E-05
LUNGS	7.87E-05	2.42E-04
THYROID	2.91E-06	9.21E-06
ENDOST	1.49E-04	4.74E-04
RMNDR	1.34E-05	4.28E-05
EFFEC	2.07E-05	6.50E-05

FREQUENCY DISTRIBUTION OF LIFETIME FATAL CANCER RISKS

Risk Range	Number of People	Number of People In This Risk Range Or Higher	Deaths/Year In This Risk Range	Deaths/Year In This Risk Range Or Higher
1.0E+00 TO 1.0E-01	0	0	0.00E+00	0.00E+00
1.0E-01 TO 1.0E-02	0	0	0.00E+00	0.00E+00
1.0E-02 TO 1.0E-03	0	0	0.00E+00	0.00E+00
1.0E-03 TO 1.0E-04	0	0	0.00E+00	0.00E+00
1.0E-04 TO 1.0E-05	0	0	0.00E+00	0.00E+00
1.0E-05 TO 1.0E-06	0	0	0.00E+00	0.00E+00
LESS THAN 1.0E-06	113910	113910	1.07E-08	1.07E-08

□

Mar 5, 2003 1:39 pm

SYNOPSIS

Page 2

RADIONUCLIDE EMISSIONS DURING THE YEAR 2002

Nuclide	Class	Size	Source #1 Ci/y	TOTAL Ci/y
AC-228	Y	1.00	1.2E-09	1.2E-09
BI-212	W	1.00	1.1E-09	1.1E-09
BI-214	W	1.00	1.9E-09	1.9E-09
C-14	*	0.00	2.0E-08	2.0E-08
CO-60	Y	1.00	2.0E-10	2.0E-10
CS-137	D	1.00	4.9E-11	4.9E-11

I.syn

PB-210	D	1.00	8.6E-09	8.6E-09
PB-212	D	1.00	1.3E-09	1.3E-09
PB-214	D	1.00	2.5E-09	2.5E-09
K-40	D	1.00	2.5E-08	2.5E-08
RA-226	W	1.00	3.3E-09	3.3E-09
SR-90	D	1.00	1.2E-09	1.2E-09
TL-208	D	1.00	4.7E-10	4.7E-10
TH-234	Y	1.00	2.8E-09	2.8E-09
U-235	Y	1.00	4.1E-10	4.1E-10
U-238	Y	1.00	2.8E-09	2.8E-09

SITE INFORMATION

Temperature: 15 degrees C
 Precipitation: 26 cm/y
 Mixing Height: 1000 m

□ Mar 5, 2003 1:39 pm

SYNOPSIS
Page 3

SOURCE INFORMATION

Source Number: 1

Source Height (m): 0.00
 Area (sq m): 5.26E+03

Plume Rise							
Pasquill Cat:	A	B	C	D	E	F	G
	<u> </u>						
Zero:	0.00	0.00	0.00	0.00	0.00	0.00	0.00

AGRICULTURAL DATA

	<u>Vegetable</u>	<u>Milk</u>	<u>Meat</u>
Fraction Home Produced:	0.700	0.399	0.442
Fraction From Assessment Area:	0.300	0.601	0.558
Fraction Imported:	0.000	0.000	0.000

Beef Cattle Density: 8.81E-02
 Milk Cattle Density: 2.85E-02
 Land Fraction Cultivated
 for Vegetable Crops: 2.50E-01

□ Mar 5, 2003 1:39 pm

SYNOPSIS
Page 4

I.syn
POPULATION DATA

Direction	Distance (m)						
	250	750	1500	2500	3500	4500	6523
N	439	439	2286	2286	4058	1191	4506
NNW	439	439	2286	4572	2624	2352	295
NW	439	439	2286	2286	2286	6417	2412
WNW	439	439	2286	2286	2286	4572	343
W	439	111	10	10	10	10	10
WSW	0	0	5	5	5	5	1
SW	0	0	2	2	2	2	2
SSW	0	0	2	2	2	2	2
S	0	0	2	2	2	2	1
SSE	0	0	2	2	2	2	0
SE	0	0	2	2	2	2	1
ESE	0	0	2	2	2	2	2
E	0	0	0	228	228	1745	2088
ENE	116	116	571	1371	1486	4681	15363
NE	439	330	115	1320	8238	4363	330
NNE	439	439	3187	3200	3187	10	13

Direction	Distance (m)						
	9023						
N	1189						
NNW	115						
NW	86						
WNW	115						
W	57						
WSW	0						
SW	0						
SSW	0						
S	0						
SSE	2						
SE	2						
ESE	2						
E	1						
ENE	460						
NE	171						
NNE	2						

B.sum

C A P 8 8 - P C

Version 1.00

Clean Air Act Assessment Package - 1988

D O S E A N D R I S K E Q U I V A L E N T S U M M A R I E S

Non-Radon Individual Assessment
Mar 7, 2003 10:26 am

Facility: LEHR
Address: Old Davis Road
City: Davis
State: CA Zip: 95616

Source Category: Nonpoint
Source Type: Area
Emission Year: 2002

Comments:

Dataset Name: EDPind
Dataset Date: Mar 7, 2003 10:26 am
Wind File: WNDFILES\SAC0320.WND

Mar 7, 2003 10:26 am

SUMMARY
Page 1

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Selected Individual (mrem/y)
GONADS	2.98E-07
BREAST	3.06E-07
R MAR	3.52E-05
LUNGS	5.33E-04

	B.sum	
THYROID		2.99E-07
ENDOST		4.36E-04
RMNDR		8.75E-07
EFFEC		8.17E-05

PATHWAY EFFECTIVE DOSE EQUIVALENT SUMMARY

Pathway	Selected Individual (mrem/y)
INGESTION	5.74E-07
INHALATION	8.11E-05
AIR IMMERSION	1.03E-10
GROUND SURFACE	1.10E-08
INTERNAL	8.17E-05
EXTERNAL	1.11E-08
TOTAL	8.17E-05

Mar 7, 2003 10:26 am

SUMMARY
Page 2

NUCLIDE EFFECTIVE DOSE EQUIVALENT SUMMARY

Nuclide	Selected Individual (mrem/y)
BI-212	1.34E-09
BI-214	7.46E-11
CS-137	1.69E-08
PB-214	6.22E-11
SR-90	4.72E-08
TL-208	3.75E-11
TH-228	3.74E-05
TH-232	4.38E-05
TH-234	6.27E-09
H-3	8.05E-11
U-235	4.78E-07
TOTAL	8.17E-05

Mar 7, 2003 10:26 am

SUMMARY
Page 3

CANCER RISK SUMMARY

Cancer	B.sum
	Selected Individual Total Lifetime Fatal Cancer Risk
LEUKEMIA	3.18E-11
BONE	2.06E-11
THYROID	5.73E-14
BREAST	4.94E-13
LUNG	9.48E-10
STOMACH	4.10E-13
BOWEL	6.24E-13
LIVER	1.74E-12
PANCREAS	2.84E-13
URINARY	2.57E-13
OTHER	3.48E-13
TOTAL	1.00E-09

PATHWAY RISK SUMMARY

Pathway	Selected Individual Total Lifetime Fatal Cancer Risk
	INGESTION
INHALATION	1.00E-09
AIR IMMERSION	2.50E-15
GROUND SURFACE	2.53E-13
INTERNAL	1.00E-09
EXTERNAL	2.56E-13
TOTAL	1.00E-09

Mar 7, 2003 10:26 am

SUMMARY
Page 4

NUCLIDE RISK SUMMARY

Nuclide	Selected Individual Total Lifetime Fatal Cancer Risk
	BI-212
BI-214	8.64E-15
CS-137	4.42E-13
PB-214	1.06E-14
SR-90	7.96E-13
TL-208	9.18E-16
TH-228	7.50E-10
TH-232	2.47E-10
TH-234	2.13E-13

	B.sum
H-3	2.18E-15
U-235	6.29E-12
TOTAL	1.00E-09

Mar 7, 2003 10:26 am

SUMMARY
Page 5

INDIVIDUAL EFFECTIVE DOSE EQUIVALENT RATE (mrem/y)
(All Radionuclides and Pathways)

Direction	Distance (m)						
	132	143	150	165	180	500	1000
N	8.2E-05	7.2E-05	6.6E-05	5.6E-05	4.7E-05	6.6E-06	1.8E-06
NNW	6.6E-05	5.5E-05	4.9E-05	4.0E-05	3.4E-05	4.7E-06	1.3E-06
NW	7.2E-05	6.2E-05	5.7E-05	4.8E-05	4.0E-05	5.6E-06	1.5E-06
WNW	5.6E-05	4.8E-05	4.4E-05	3.6E-05	3.1E-05	4.2E-06	1.1E-06
W	3.9E-05	3.3E-05	3.0E-05	2.5E-05	2.1E-05	2.9E-06	7.8E-07
WSW	1.8E-05	1.5E-05	1.3E-05	1.0E-05	8.7E-06	1.2E-06	3.5E-07
SW	1.1E-05	9.3E-06	8.4E-06	6.9E-06	5.8E-06	8.3E-07	2.5E-07
SSW	1.2E-05	1.0E-05	8.9E-06	7.1E-06	5.9E-06	8.5E-07	2.6E-07
S	2.9E-05	2.5E-05	2.3E-05	1.9E-05	1.6E-05	2.3E-06	6.3E-07
SSE	3.7E-05	3.3E-05	3.0E-05	2.6E-05	2.2E-05	3.1E-06	8.5E-07
SE	2.4E-05	2.0E-05	1.8E-05	1.5E-05	1.2E-05	1.8E-06	5.1E-07
ESE	1.4E-05	1.2E-05	1.0E-05	8.5E-06	7.1E-06	1.0E-06	3.1E-07
E	1.5E-05	1.3E-05	1.1E-05	9.5E-06	8.0E-06	1.1E-06	3.3E-07
ENE	1.8E-05	1.5E-05	1.3E-05	1.0E-05	8.7E-06	1.3E-06	3.7E-07
NE	4.3E-05	3.7E-05	3.4E-05	2.9E-05	2.4E-05	3.4E-06	9.4E-07
NNE	5.5E-05	4.6E-05	4.2E-05	3.4E-05	2.9E-05	4.1E-06	1.1E-06

Direction	Distance (m)
	187
N	4.4E-05
NNW	3.2E-05
NW	3.8E-05
WNW	2.8E-05
W	2.0E-05
WSW	8.1E-06
SW	5.4E-06
SSW	5.5E-06
S	1.5E-05
SSE	2.0E-05
SE	1.2E-05
ESE	6.6E-06
E	7.4E-06
ENE	8.1E-06
NE	2.3E-05
NNE	2.7E-05

B.sum

Mar 7, 2003 10:26 am

SUMMARY
Page 6

INDIVIDUAL LIFETIME RISK (deaths)
(All Radionuclides and Pathways)

Direction	Distance (m)						
	132	143	150	165	180	500	1000
N	1.0E-09	8.8E-10	8.1E-10	6.9E-10	5.8E-10	8.1E-11	2.1E-11
NNW	8.1E-10	6.8E-10	6.1E-10	5.0E-10	4.2E-10	5.8E-11	1.5E-11
NW	8.8E-10	7.7E-10	7.0E-10	5.9E-10	5.0E-10	6.9E-11	1.8E-11
WNW	6.9E-10	5.9E-10	5.4E-10	4.5E-10	3.8E-10	5.1E-11	1.3E-11
W	4.7E-10	4.1E-10	3.7E-10	3.1E-10	2.6E-10	3.5E-11	9.3E-12
WSW	2.2E-10	1.8E-10	1.6E-10	1.3E-10	1.1E-10	1.5E-11	3.9E-12
SW	1.4E-10	1.1E-10	1.0E-10	8.4E-11	7.1E-11	9.8E-12	2.8E-12
SSW	1.5E-10	1.2E-10	1.1E-10	8.7E-11	7.3E-11	1.0E-11	2.8E-12
S	3.5E-10	3.1E-10	2.8E-10	2.3E-10	2.0E-10	2.7E-11	7.4E-12
SSE	4.6E-10	4.0E-10	3.7E-10	3.2E-10	2.7E-10	3.8E-11	1.0E-11
SE	2.9E-10	2.4E-10	2.2E-10	1.8E-10	1.5E-10	2.1E-11	5.9E-12
ESE	1.7E-10	1.4E-10	1.3E-10	1.0E-10	8.7E-11	1.2E-11	3.4E-12
E	1.8E-10	1.5E-10	1.4E-10	1.2E-10	9.8E-11	1.4E-11	3.8E-12
ENE	2.3E-10	1.8E-10	1.6E-10	1.3E-10	1.1E-10	1.5E-11	4.2E-12
NE	5.3E-10	4.6E-10	4.2E-10	3.5E-10	3.0E-10	4.2E-11	1.1E-11
NNE	6.7E-10	5.7E-10	5.1E-10	4.2E-10	3.5E-10	5.0E-11	1.3E-11

Direction	Distance (m)
	187
N	5.4E-10
NNW	3.9E-10
NW	4.6E-10
WNW	3.5E-10
W	2.4E-10
WSW	9.9E-11
SW	6.6E-11
SSW	6.7E-11
S	1.8E-10
SSE	2.5E-10
SE	1.4E-10
ESE	8.1E-11
E	9.1E-11
ENE	1.0E-10
NE	2.8E-10
NNE	3.3E-10

C.syn

C A P 8 8 - P C

Version 1.00

Clean Air Act Assessment Package - 1988

S Y N O P S I S R E P O R T

Non-Radon Population Assessment
Mar 7, 2003 10:26 am

Facility: LEHR
Address: Old Davis Road
City: Davis
State: CA Zip: 95616

Effective Dose Equivalent
(mrem/year)

2.67E-05

At This Location: 250 Meters North
Source Category: Nonpoint
Source Type: Area
Emission Year: 2002

Comments:

Dataset Name: EDppop
Dataset Date: Mar 7, 2003 10:24 am
Wind File: WNDFILES\SAC0320.WND
Population File: POPFILES\02LEHR.POP

Mar 7, 2003 10:26 am

SYNOPSIS
Page 1

MAXIMALLY EXPOSED INDIVIDUAL

Page 1

C.syn

Location Of The Individual: 250 Meters North
Lifetime Fatal Cancer Risk: 3.41E-10

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Selected Individual (mrem/y)	Collective Population (person-rem/y)
GONADS	9.36E-08	2.89E-07
BREAST	9.60E-08	2.96E-07
R MAR	1.10E-05	3.37E-05
LUNGS	1.77E-04	5.43E-04
THYROID	9.36E-08	2.89E-07
ENDOST	1.36E-04	4.17E-04
RMNDR	2.79E-07	8.68E-07
EFFEC	2.67E-05	8.21E-05

FREQUENCY DISTRIBUTION OF LIFETIME FATAL CANCER RISKS

Risk Range	Number of People	Number of People In This Risk Range Or Higher	Deaths/Year In This Risk Range	Deaths/Year In This Risk Range Or Higher
1.0E+00 TO 1.0E-01	0	0	0.00E+00	0.00E+00
1.0E-01 TO 1.0E-02	0	0	0.00E+00	0.00E+00
1.0E-02 TO 1.0E-03	0	0	0.00E+00	0.00E+00
1.0E-03 TO 1.0E-04	0	0	0.00E+00	0.00E+00
1.0E-04 TO 1.0E-05	0	0	0.00E+00	0.00E+00
1.0E-05 TO 1.0E-06	0	0	0.00E+00	0.00E+00
LESS THAN 1.0E-06	113910	113910	1.48E-08	1.48E-08

□

Mar 7, 2003 10:26 am

SYNOPSIS
Page 2

RADIONUCLIDE EMISSIONS DURING THE YEAR 2002

Nuclide	Class	Size	Source #1 Ci/y	TOTAL Ci/y
BI-212	W	1.00	5.4E-10	5.4E-10
BI-214	W	1.00	7.4E-10	7.4E-10
CS-137	D	1.00	2.5E-10	2.5E-10
SR-90	D	1.00	2.1E-10	2.1E-10
TL-208	D	1.00	2.8E-10	2.8E-10
TH-232	Y	1.00	1.6E-09	1.6E-09

				C.syn
H-3	*	0.00	1.6E-09	1.6E-09
U-235	Y	1.00	5.0E-11	5.0E-11
TH-228	Y	1.00	2.3E-09	2.3E-09
TH-234	Y	1.00	1.1E-09	1.1E-09
PB-214	D	1.00	7.9E-10	7.9E-10

SITE INFORMATION

Temperature: 15 degrees C
 Precipitation: 26 cm/y
 Mixing Height: 1000 m

□ Mar 7, 2003 10:26 am

SYNOPSIS
 Page 3

SOURCE INFORMATION

Source Number: 1

Source Height (m): 0.00
 Area (sq m): 3.90E+03

Plume Rise							
Pasquill Cat:	A	B	C	D	E	F	G
Zero:	0.00	0.00	0.00	0.00	0.00	0.00	0.00

AGRICULTURAL DATA

	Vegetable	Milk	Meat
Fraction Home Produced:	0.700	0.399	0.442
Fraction From Assessment Area:	0.300	0.601	0.558
Fraction Imported:	0.000	0.000	0.000
Beef Cattle Density:	8.81E-02		
Milk Cattle Density:	2.85E-02		
Land Fraction Cultivated for Vegetable Crops:	2.50E-01		

□ Mar 7, 2003 10:26 am

SYNOPSIS
 Page 4

POPULATION DATA

Distance (m)

Direction	C.syn						
	250	750	1500	2500	3500	4500	6523
N	439	439	2286	2286	4058	1191	4506
NNW	439	439	2286	4572	2624	2352	295
NW	439	439	2286	2286	2286	6417	2412
WNW	439	439	2286	2286	2286	4572	343
W	439	111	10	10	10	10	10
WSW	0	0	5	5	5	5	1
SW	0	0	2	2	2	2	2
SSW	0	0	2	2	2	2	2
S	0	0	2	2	2	2	1
SSE	0	0	2	2	2	2	0
SE	0	0	2	2	2	2	1
ESE	0	0	2	2	2	2	2
E	0	0	0	228	228	1745	2088
ENE	116	116	571	1371	1486	4681	15363
NE	439	330	115	1320	8238	4363	330
NNE	439	439	3187	3200	3187	10	13

Direction	Distance (m)						
	9023						
N	1189						
NNW	115						
NW	86						
WNW	115						
W	57						
WSW	0						
SW	0						
SSW	0						
S	0						
SSE	2						
SE	2						
ESE	2						
E	1						
ENE	460						
NE	171						
NNE	2						

D.sum

C A P 8 8 - P C

Version 1.00

Clean Air Act Assessment Package - 1988

D O S E A N D R I S K E Q U I V A L E N T S U M M A R I E S

Non-Radon Individual Assessment

Mar 31, 2003 1:33 pm

Facility: LEHR
Address: Old Davis Road
City: Davis
State: CA Zip: 95616

Source Category: Non-point
Source Type: Area
Emission Year: 2002

Comments:

Dataset Name: DSS3ind
Dataset Date: Mar 31, 2003 1:11 pm
Wind File: WNDFILES\SAC0320.WND

Mar 31, 2003 1:33 pm

SUMMARY
Page 1

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Selected Individual (mrem/y)
GONADS	1.37E-06
BREAST	1.33E-06
R MAR	1.49E-05
LUNGS	1.69E-04

	D.sum	
THYROID		1.31E-06
ENDOST		1.76E-04
RMNDR		1.23E-05
EFFEC		3.17E-05

PATHWAY EFFECTIVE DOSE EQUIVALENT SUMMARY

Pathway	Selected Individual (mrem/y)
INGESTION	8.41E-06
INHALATION	2.32E-05
AIR IMMERSION	2.05E-10
GROUND SURFACE	4.49E-08
INTERNAL	3.16E-05
EXTERNAL	4.51E-08
TOTAL	3.17E-05

□ Mar 31, 2003 1:33 pm

SUMMARY
Page 2

NUCLIDE EFFECTIVE DOSE EQUIVALENT SUMMARY

Nuclide	Selected Individual (mrem/y)
BI-214	2.51E-10
C-14	4.14E-10
CS-137	4.06E-09
PB-210	7.96E-06
PB-214	2.08E-10
PU-241	3.01E-07
RA-226	3.25E-06
SR-90	4.51E-07
TH-234	4.95E-09
U-233	1.24E-05
U-235	9.45E-07
U-238	6.39E-06
TOTAL	3.17E-05

□ Mar 31, 2003 1:33 pm

SUMMARY
Page 3

CANCER RISK SUMMARY

D.sum

Cancer	Selected Individual Total Lifetime Fatal Cancer Risk
LEUKEMIA	2.00E-11
BONE	9.71E-12
THYROID	2.24E-13
BREAST	1.89E-12
LUNG	2.90E-10
STOMACH	1.44E-12
BOWEL	1.14E-12
LIVER	3.33E-11
PANCREAS	1.08E-12
URINARY	8.37E-12
OTHER	1.32E-12
TOTAL	3.69E-10

PATHWAY RISK SUMMARY

Pathway	Selected Individual Total Lifetime Fatal Cancer Risk
INGESTION	5.91E-11
INHALATION	3.09E-10
AIR IMMERSION	4.94E-15
GROUND SURFACE	1.04E-12
INTERNAL	3.68E-10
EXTERNAL	1.04E-12
TOTAL	3.69E-10

□

Mar 31, 2003 1:33 pm

SUMMARY
Page 4

NUCLIDE RISK SUMMARY

Nuclide	Selected Individual Total Lifetime Fatal Cancer Risk
BI-214	2.91E-14
C-14	1.01E-14
CS-137	1.06E-13
PB-210	5.67E-11
PB-214	3.56E-14
PU-241	1.15E-12
RA-226	4.68E-11
SR-90	7.60E-12

	D. sum
TH-234	1.72E-13
U-233	1.60E-10
U-235	1.25E-11
U-238	8.35E-11
TOTAL	3.69E-10

□

Mar 31, 2003 1:33 pm

SUMMARY
Page 5

INDIVIDUAL EFFECTIVE DOSE EQUIVALENT RATE (mrem/y)
(All Radionuclides and Pathways)

Direction	Distance (m)						
	15	24	34	53	85	110	118
N	3.0E-05	1.3E-05	7.3E-06	3.7E-06	1.8E-06	1.2E-06	1.1E-06
NNW	3.2E-05	1.4E-05	7.1E-06	3.0E-06	1.4E-06	9.8E-07	9.0E-07
NW	2.9E-05	1.2E-05	6.8E-06	3.3E-06	1.6E-06	1.1E-06	1.0E-06
WNW	2.4E-05	1.1E-05	5.8E-06	2.7E-06	1.3E-06	9.2E-07	8.5E-07
W	1.8E-05	7.1E-06	4.0E-06	2.0E-06	1.0E-06	7.5E-07	7.0E-07
WSW	1.2E-05	4.3E-06	2.3E-06	1.1E-06	6.3E-07	5.2E-07	5.0E-07
SW	9.0E-06	2.6E-06	1.5E-06	8.2E-07	5.4E-07	4.7E-07	4.5E-07
SSW	9.6E-06	3.4E-06	1.8E-06	8.8E-07	5.4E-07	4.7E-07	4.6E-07
S	1.1E-05	5.2E-06	3.0E-06	1.5E-06	8.4E-07	6.5E-07	6.2E-07
SSE	1.2E-05	6.0E-06	3.5E-06	1.9E-06	1.0E-06	7.6E-07	7.1E-07
SE	1.2E-05	4.9E-06	2.7E-06	1.3E-06	7.4E-07	5.9E-07	5.6E-07
ESE	9.8E-06	3.4E-06	1.9E-06	9.4E-07	5.8E-07	4.9E-07	4.7E-07
E	9.9E-06	3.0E-06	1.8E-06	9.7E-07	6.0E-07	5.1E-07	4.9E-07
ENE	1.3E-05	4.8E-06	2.5E-06	1.1E-06	6.2E-07	5.2E-07	5.0E-07
NE	2.0E-05	7.2E-06	4.1E-06	2.1E-06	1.1E-06	8.0E-07	7.5E-07
NNE	2.6E-05	1.1E-05	5.9E-06	2.6E-06	1.2E-06	8.8E-07	8.2E-07

Direction	Distance (m)		
	346	354	1173
N	4.6E-07	4.5E-07	3.7E-07
NNW	4.3E-07	4.3E-07	3.7E-07
NW	4.4E-07	4.4E-07	3.7E-07
WNW	4.2E-07	4.2E-07	3.7E-07
W	4.0E-07	4.0E-07	3.7E-07
WSW	3.8E-07	3.8E-07	3.6E-07
SW	3.7E-07	3.7E-07	3.6E-07
SSW	3.7E-07	3.7E-07	3.6E-07
S	3.9E-07	3.9E-07	3.6E-07
SSE	4.1E-07	4.0E-07	3.7E-07
SE	3.9E-07	3.9E-07	3.6E-07
ESE	3.8E-07	3.7E-07	3.6E-07
E	3.8E-07	3.8E-07	3.6E-07
ENE	3.8E-07	3.8E-07	3.6E-07

			D.sum
NE	4.1E-07	4.1E-07	3.7E-07
NNE	4.2E-07	4.2E-07	3.7E-07

□ Mar 31, 2003 1:33 pm

SUMMARY
Page 6

INDIVIDUAL LIFETIME RISK (deaths)
(All Radionuclides and Pathways)

	Distance (m)						
Direction	15	24	34	53	85	110	118
N	3.5E-10	1.5E-10	8.4E-11	4.1E-11	1.9E-11	1.3E-11	1.1E-11
NNW	3.7E-10	1.6E-10	8.2E-11	3.4E-11	1.4E-11	9.8E-12	8.9E-12
NW	3.4E-10	1.4E-10	7.8E-11	3.7E-11	1.7E-11	1.1E-11	1.0E-11
WNW	2.7E-10	1.2E-10	6.6E-11	3.0E-11	1.3E-11	9.2E-12	8.3E-12
W	2.1E-10	8.1E-11	4.5E-11	2.1E-11	1.0E-11	7.1E-12	6.6E-12
WSW	1.4E-10	4.8E-11	2.6E-11	1.1E-11	5.7E-12	4.5E-12	4.2E-12
SW	1.0E-10	2.9E-11	1.6E-11	8.0E-12	4.6E-12	3.8E-12	3.7E-12
SSW	1.1E-10	3.8E-11	2.0E-11	8.7E-12	4.7E-12	3.9E-12	3.7E-12
S	1.3E-10	5.9E-11	3.3E-11	1.6E-11	8.2E-12	6.0E-12	5.6E-12
SSE	1.4E-10	6.8E-11	4.0E-11	2.0E-11	1.0E-11	7.2E-12	6.6E-12
SE	1.3E-10	5.6E-11	3.0E-11	1.4E-11	7.0E-12	5.2E-12	4.9E-12
ESE	1.1E-10	3.8E-11	2.0E-11	9.4E-12	5.1E-12	4.1E-12	3.9E-12
E	1.1E-10	3.4E-11	1.9E-11	9.7E-12	5.4E-12	4.3E-12	4.1E-12
ENE	1.5E-10	5.5E-11	2.8E-11	1.1E-11	5.6E-12	4.4E-12	4.2E-12
NE	2.3E-10	8.3E-11	4.7E-11	2.3E-11	1.1E-11	7.7E-12	7.1E-12
NNE	3.0E-10	1.3E-10	6.8E-11	2.9E-11	1.3E-11	8.7E-12	7.9E-12

	Distance (m)		
Direction	346	354	1173
N	3.7E-12	3.6E-12	2.7E-12
NNW	3.4E-12	3.3E-12	2.7E-12
NW	3.5E-12	3.5E-12	2.7E-12
WNW	3.3E-12	3.3E-12	2.7E-12
W	3.1E-12	3.0E-12	2.6E-12
WSW	2.8E-12	2.8E-12	2.6E-12
SW	2.7E-12	2.7E-12	2.6E-12
SSW	2.7E-12	2.7E-12	2.6E-12
S	3.0E-12	2.9E-12	2.6E-12
SSE	3.1E-12	3.1E-12	2.6E-12
SE	2.9E-12	2.9E-12	2.6E-12
ESE	2.7E-12	2.7E-12	2.6E-12
E	2.8E-12	2.8E-12	2.6E-12
ENE	2.8E-12	2.8E-12	2.6E-12
NE	3.2E-12	3.1E-12	2.6E-12

			D.sum
NNE	3.3E-12	3.2E-12	2.7E-12

E.syn

C A P 8 8 - P C

Version 1.00

Clean Air Act Assessment Package - 1988

S Y N O P S I S R E P O R T

Non-Radon Population Assessment
Mar 31, 2003 1:34 pm

Facility: LEHR
Address: Old Davis Road
City: Davis
State: CA Zip: 95616

Effective Dose Equivalent
(mrem/year)

1.76E-07

At This Location: 250 Meters North
Source Category: Non-point
Source Type: Area
Emission Year: 2002

Comments:

Dataset Name: DSS3pop
Dataset Date: Mar 31, 2003 1:13 pm
Wind File: WNDFILES\SAC0320.WND
Population File: POPFILES\02LEHR.POP

Mar 31, 2003 1:34 pm

SYNOPSIS
Page 1

MAXIMALLY EXPOSED INDIVIDUAL

Page 1

E.syn

Location Of The Individual: 250 Meters North
Lifetime Fatal Cancer Risk: 2.06E-12

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Selected Individual (mrem/y)	Collective Population (person-rem/y)
GONADS	7.41E-09	2.36E-08
BREAST	7.19E-09	2.29E-08
R MAR	7.95E-08	2.54E-07
LUNGS	9.56E-07	2.94E-06
THYROID	7.09E-09	2.26E-08
ENDOST	9.46E-07	3.02E-06
RMNDR	6.63E-08	2.12E-07
EFFEC	1.76E-07	5.47E-07

FREQUENCY DISTRIBUTION OF LIFETIME FATAL CANCER RISKS

Risk Range	Number of People	Number of People In This Risk Range Or Higher	Deaths/Year In This Risk Range	Deaths/Year In This Risk Range Or Higher
1.0E+00 TO 1.0E-01	0	0	0.00E+00	0.00E+00
1.0E-01 TO 1.0E-02	0	0	0.00E+00	0.00E+00
1.0E-02 TO 1.0E-03	0	0	0.00E+00	0.00E+00
1.0E-03 TO 1.0E-04	0	0	0.00E+00	0.00E+00
1.0E-04 TO 1.0E-05	0	0	0.00E+00	0.00E+00
1.0E-05 TO 1.0E-06	0	0	0.00E+00	0.00E+00
LESS THAN 1.0E-06	113910	113910	9.01E-11	9.01E-11

□

Mar 31, 2003 1:34 pm

SYNOPSIS
Page 2

RADIONUCLIDE EMISSIONS DURING THE YEAR 2002

Nuclide	Class	Size	Source #1 Ci/y	TOTAL Ci/y
BI-214	W	1.00	4.4E-11	4.4E-11
C-14	*	0.00	3.1E-12	3.1E-12
CS-137	D	1.00	1.2E-12	1.2E-12
PB-210	D	1.00	4.3E-11	4.3E-11
PB-214	D	1.00	4.7E-11	4.7E-11
PU-241	Y	1.00	1.3E-11	1.3E-11

				E.syn
RA-226	W	1.00	4.9E-11	4.9E-11
SR-90	D	1.00	4.0E-11	4.0E-11
TH-234	Y	1.00	1.7E-11	1.7E-11
U-233	Y	1.00	2.2E-11	2.2E-11
U-235	Y	1.00	1.8E-12	1.8E-12
U-238	Y	1.00	1.3E-11	1.3E-11

SITE INFORMATION

Temperature: 15 degrees C
 Precipitation: 26 cm/y
 Mixing Height: 1000 m

□ Mar 31, 2003 1:34 pm

SYNOPSIS
 Page 3

SOURCE INFORMATION

Source Number: 1

Source Height (m): 0.00
 Area (sq m): 6.52E+02

Plume Rise							
Pasquill Cat:	A	B	C	D	E	F	G
Zero:	0.00	0.00	0.00	0.00	0.00	0.00	0.00

AGRICULTURAL DATA

	Vegetable	Milk	Meat
Fraction Home Produced:	0.700	0.399	0.442
Fraction From Assessment Area:	0.300	0.601	0.558
Fraction Imported:	0.000	0.000	0.000

Beef Cattle Density: 8.81E-02
 Milk Cattle Density: 2.85E-02
 Land Fraction Cultivated
 for Vegetable Crops: 2.50E-01

□ Mar 31, 2003 1:34 pm

SYNOPSIS
 Page 4

POPULATION DATA

Distance (m)

E. syn

Direction	250	750	1500	2500	3500	4500	6523
N	439	439	2286	2286	4058	1191	4506
NNW	439	439	2286	4572	2624	2352	295
NW	439	439	2286	2286	2286	6417	2412
WNW	439	439	2286	2286	2286	4572	343
W	439	111	10	10	10	10	10
WSW	0	0	5	5	5	5	1
SW	0	0	2	2	2	2	2
SSW	0	0	2	2	2	2	2
S	0	0	2	2	2	2	1
SSE	0	0	2	2	2	2	0
SE	0	0	2	2	2	2	1
ESE	0	0	2	2	2	2	2
E	0	0	0	228	228	1745	2088
ENE	116	116	571	1371	1486	4681	15363
NE	439	330	115	1320	8238	4363	330
NNE	439	439	3187	3200	3187	10	13

Distance (m)

Direction	9023
N	1189
NNW	115
NW	86
WNW	115
W	57
WSW	0
SW	0
SSW	0
S	0
SSE	2
SE	2
ESE	2
E	1
ENE	460
NE	171
NNE	2

G.syn

C A P 8 8 - P C

Version 1.00

Clean Air Act Assessment Package - 1988

S Y N O P S I S R E P O R T

Non-Radon Population Assessment
Mar 31, 2003 1:38 pm

Facility: LEHR
Address: Old Davis Road
City: Davis
State: CA Zip: 95616

Effective Dose Equivalent
(mrem/year)

1.70E-07

At This Location: 250 Meters North
Source Category: nonpoint
Source Type: Area
Emission Year: 2002

Comments:

Dataset Name: DSS6pop
Dataset Date: Mar 31, 2003 1:21 pm
Wind File: WNDFILES\SAC0320.WND
Population File: POPFILES\02LEHR.POP

Mar 31, 2003 1:38 pm

SYNOPSIS
Page 1

MAXIMALLY EXPOSED INDIVIDUAL

Page 1

G.syn

Location Of The Individual: 250 Meters North
Lifetime Fatal Cancer Risk: 1.75E-12

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Selected Individual (mrem/y)	Collective Population (person-rem/y)
GONADS	8.33E-09	2.63E-08
BREAST	6.61E-09	2.10E-08
R MAR	8.93E-08	2.84E-07
LUNGS	7.51E-07	2.31E-06
THYROID	6.51E-09	2.07E-08
ENDOST	1.21E-06	3.85E-06
RMNDR	9.73E-08	3.10E-07
EFFEC	1.70E-07	5.30E-07

FREQUENCY DISTRIBUTION OF LIFETIME FATAL CANCER RISKS

Risk Range	Number of People	Number of People In This Risk Range Or Higher	Deaths/Year In This Risk Range	Deaths/Year In This Risk Range Or Higher
1.0E+00 TO 1.0E-01	0	0	0.00E+00	0.00E+00
1.0E-01 TO 1.0E-02	0	0	0.00E+00	0.00E+00
1.0E-02 TO 1.0E-03	0	0	0.00E+00	0.00E+00
1.0E-03 TO 1.0E-04	0	0	0.00E+00	0.00E+00
1.0E-04 TO 1.0E-05	0	0	0.00E+00	0.00E+00
1.0E-05 TO 1.0E-06	0	0	0.00E+00	0.00E+00
LESS THAN 1.0E-06	113910	113910	7.67E-11	7.67E-11

□

Mar 31, 2003 1:38 pm

SYNOPSIS
Page 2

RADIONUCLIDE EMISSIONS DURING THE YEAR 2002

Nuclide	Class	Size	Source #1 Ci/y	TOTAL Ci/y
AC-228	Y	1.00	2.6E-11	2.6E-11
AM-241	W	1.00	7.0E-13	7.0E-13
BI-212	W	1.00	1.7E-11	1.7E-11
BI-214	W	1.00	2.3E-11	2.3E-11
CS-137	D	1.00	2.1E-12	2.1E-12
CO-60	Y	1.00	5.0E-13	5.0E-13

G.syn

PB-210	D	1.00	6.7E-11	6.7E-11
PB-214	D	1.00	2.9E-11	2.9E-11
SR-90	D	1.00	8.1E-12	8.1E-12
TL-208	D	1.00	9.2E-12	9.2E-12
H-3	*	0.00	8.8E-11	8.8E-11
U-233	Y	1.00	2.8E-11	2.8E-11
U-235	Y	1.00	1.7E-12	1.7E-12

SITE INFORMATION

Temperature: 15 degrees C
 Precipitation: 26 cm/y
 Mixing Height: 1000 m

□ Mar 31, 2003 1:38 pm

SYNOPSIS
Page 3

SOURCE INFORMATION

Source Number: 1

Source Height (m): 0.00
 Area (sq m): 1.24E+03

Plume Rise							
Pasquill Cat:	A	B	C	D	E	F	G
<hr/>							
Zero:	0.00	0.00	0.00	0.00	0.00	0.00	0.00

AGRICULTURAL DATA

	<u>Vegetable</u>	<u>Milk</u>	<u>Meat</u>
Fraction Home Produced:	0.700	0.399	0.442
Fraction From Assessment Area:	0.300	0.601	0.558
Fraction Imported:	0.000	0.000	0.000

Beef Cattle Density: 8.81E-02
 Milk Cattle Density: 2.85E-02
 Land Fraction Cultivated
 for Vegetable Crops: 2.50E-01

□ Mar 31, 2003 1:38 pm

SYNOPSIS
Page 4

POPULATION DATA

G.syn
Distance (m)

Direction	250	750	1500	2500	3500	4500	6523
N	439	439	2286	2286	4058	1191	4506
NNW	439	439	2286	4572	2624	2352	295
NW	439	439	2286	2286	2286	6417	2412
WNW	439	439	2286	2286	2286	4572	343
W	439	111	10	10	10	10	10
WSW	0	0	5	5	5	5	1
SW	0	0	2	2	2	2	2
SSW	0	0	2	2	2	2	2
S	0	0	2	2	2	2	1
SSE	0	0	2	2	2	2	0
SE	0	0	2	2	2	2	1
ESE	0	0	2	2	2	2	2
E	0	0	0	228	228	1745	2088
ENE	116	116	571	1371	1486	4681	15363
NE	439	330	115	1320	8238	4363	330
NNE	439	439	3187	3200	3187	10	13

Distance (m)

Direction	9023
N	1189
NNW	115
NW	86
WNW	115
W	57
WSW	0
SW	0
SSW	0
S	0
SSE	2
SE	2
ESE	2
E	1
ENE	460
NE	171
NNE	2

F.sum

C A P 8 8 - P C

Version 1.00

Clean Air Act Assessment Package - 1988

D O S E A N D R I S K E Q U I V A L E N T S U M M A R I E S

Non-Radon Individual Assessment
Mar 31, 2003 1:37 pm

Facility: LEHR
Address: Old Davis Road
City: Davis
State: CA Zip: 95616

Source Category: nonpoint
Source Type: Area
Emission Year: 2002

Comments:

Dataset Name: DSS6ind
Dataset Date: Mar 31, 2003 1:20 pm
Wind File: WNDFILES\SAC0320.WND

Mar 31, 2003 1:37 pm

SUMMARY
Page 1

ORGAN DOSE EQUIVALENT SUMMARY

Organ	Selected Individual (mrem/y)
GONADS	4.25E-07
BREAST	3.41E-07
R MAR	4.60E-06
LUNGS	3.65E-05

	F.sum
THYROID	3.36E-07
ENDOST	6.23E-05
RMNDR	5.01E-06
EFFEC	8.48E-06

PATHWAY EFFECTIVE DOSE EQUIVALENT SUMMARY

Pathway	Selected Individual (mrem/y)
INGESTION	2.69E-06
INHALATION	5.78E-06
AIR IMMERSION	7.30E-11
GROUND SURFACE	9.70E-09
INTERNAL	8.47E-06
EXTERNAL	9.77E-09
TOTAL	8.48E-06

□ Mar 31, 2003 1:37 pm

SUMMARY
Page 2

NUCLIDE EFFECTIVE DOSE EQUIVALENT SUMMARY

Nuclide	Selected Individual (mrem/y)
AC-228	2.46E-09
AM-241	4.37E-07
BI-212	6.50E-10
BI-214	3.64E-11
CS-137	1.98E-09
CO-60	3.17E-09
PB-210	3.44E-06
PB-214	3.48E-11
SR-90	2.55E-08
TL-208	2.31E-11
H-3	6.25E-11
U-233	4.32E-06
U-235	2.46E-07
TOTAL	8.48E-06

□ Mar 31, 2003 1:37 pm

SUMMARY
Page 3

CANCER RISK SUMMARY

F.sum

Cancer	Selected Individual Total Lifetime Fatal Cancer Risk
LEUKEMIA	5.12E-12
BONE	3.28E-12
THYROID	5.15E-14
BREAST	4.21E-13
LUNG	5.87E-11
STOMACH	3.19E-13
BOWEL	1.87E-13
LIVER	1.50E-11
PANCREAS	2.40E-13
URINARY	2.96E-12
OTHER	2.93E-13
TOTAL	8.66E-11

PATHWAY RISK SUMMARY

Pathway	Selected Individual Total Lifetime Fatal Cancer Risk
INGESTION	1.89E-11
INHALATION	6.74E-11
AIR IMMERSION	1.77E-15
GROUND SURFACE	2.23E-13
INTERNAL	8.64E-11
EXTERNAL	2.25E-13
TOTAL	8.66E-11

□

Mar 31, 2003 1:37 pm

SUMMARY
Page 4

NUCLIDE RISK SUMMARY

Nuclide	Selected Individual Total Lifetime Fatal Cancer Risk
AC-228	4.96E-14
AM-241	2.23E-12
BI-212	8.35E-15
BI-214	4.22E-15
CS-137	5.18E-14
CO-60	7.95E-14
PB-210	2.45E-11

F.sum

PB-214	5.95E-15
SR-90	4.29E-13
TL-208	5.65E-16
H-3	1.69E-15
U-233	5.59E-11
U-235	3.24E-12
TOTAL	8.66E-11

Mar 31, 2003 1:37 pm

SUMMARY
Page 5

INDIVIDUAL EFFECTIVE DOSE EQUIVALENT RATE (mrem/y)
(All Radionuclides and Pathways)

Direction	Distance (m)						
	31	43	52	58	61	338	360
N	7.7E-06	4.4E-06	3.2E-06	2.7E-06	2.5E-06	2.7E-07	2.6E-07
NNW	8.5E-06	4.5E-06	3.1E-06	2.5E-06	2.2E-06	2.5E-07	2.4E-07
NW	7.4E-06	4.2E-06	3.0E-06	2.5E-06	2.3E-06	2.6E-07	2.5E-07
WNW	6.5E-06	3.6E-06	2.5E-06	2.1E-06	1.9E-06	2.4E-07	2.3E-07
W	4.4E-06	2.5E-06	1.8E-06	1.5E-06	1.3E-06	2.2E-07	2.2E-07
WSW	2.7E-06	1.4E-06	1.0E-06	8.3E-07	7.6E-07	2.0E-07	1.9E-07
SW	1.6E-06	9.2E-07	6.7E-07	5.7E-07	5.3E-07	1.9E-07	1.9E-07
SSW	2.1E-06	1.1E-06	7.9E-07	6.5E-07	6.0E-07	1.9E-07	1.9E-07
S	3.2E-06	1.8E-06	1.3E-06	1.1E-06	1.0E-06	2.1E-07	2.1E-07
SSE	3.6E-06	2.1E-06	1.6E-06	1.3E-06	1.2E-06	2.2E-07	2.2E-07
SE	3.0E-06	1.7E-06	1.2E-06	9.8E-07	9.0E-07	2.0E-07	2.0E-07
ESE	2.1E-06	1.1E-06	8.2E-07	6.9E-07	6.3E-07	1.9E-07	1.9E-07
E	1.8E-06	1.1E-06	7.9E-07	6.7E-07	6.3E-07	2.0E-07	1.9E-07
ENE	3.0E-06	1.6E-06	1.1E-06	8.7E-07	7.9E-07	2.0E-07	1.9E-07
NE	4.4E-06	2.5E-06	1.8E-06	1.5E-06	1.4E-06	2.3E-07	2.2E-07
NNE	7.0E-06	3.7E-06	2.6E-06	2.1E-06	1.9E-06	2.4E-07	2.3E-07

Direction	Distance (m)
	1246
N	1.9E-07
NNW	1.8E-07
NW	1.9E-07
WNW	1.8E-07
W	1.8E-07
WSW	1.8E-07
SW	1.8E-07
SSW	1.8E-07
S	1.8E-07
SSE	1.8E-07
SE	1.8E-07
ESE	1.8E-07

F.sum

E 1.8E-07
ENE 1.8E-07
NE 1.8E-07
NNE 1.8E-07

□

Mar 31, 2003 1:37 pm

SUMMARY
Page 6

INDIVIDUAL LIFETIME RISK (deaths)
(All Radionuclides and Pathways)

Direction	Distance (m)						
	31	43	52	58	61	338	360
N	7.9E-11	4.5E-11	3.3E-11	2.7E-11	2.5E-11	2.2E-12	2.1E-12
NNW	8.7E-11	4.5E-11	3.1E-11	2.5E-11	2.2E-11	2.0E-12	1.9E-12
NW	7.6E-11	4.2E-11	3.0E-11	2.5E-11	2.3E-11	2.1E-12	2.0E-12
WNW	6.7E-11	3.6E-11	2.5E-11	2.1E-11	1.9E-11	1.9E-12	1.8E-12
W	4.4E-11	2.5E-11	1.7E-11	1.4E-11	1.3E-11	1.7E-12	1.6E-12
WSW	2.7E-11	1.4E-11	9.8E-12	7.9E-12	7.2E-12	1.4E-12	1.4E-12
SW	1.6E-11	8.8E-12	6.4E-12	5.3E-12	4.9E-12	1.4E-12	1.4E-12
SSW	2.1E-11	1.1E-11	7.6E-12	6.2E-12	5.6E-12	1.4E-12	1.4E-12
S	3.2E-11	1.8E-11	1.3E-11	1.1E-11	1.0E-11	1.6E-12	1.6E-12
SSE	3.7E-11	2.1E-11	1.6E-11	1.3E-11	1.2E-11	1.7E-12	1.7E-12
SE	3.0E-11	1.6E-11	1.2E-11	9.5E-12	8.7E-12	1.5E-12	1.5E-12
ESE	2.1E-11	1.1E-11	7.9E-12	6.5E-12	5.9E-12	1.4E-12	1.4E-12
E	1.8E-11	1.0E-11	7.5E-12	6.3E-12	5.9E-12	1.4E-12	1.4E-12
ENE	3.0E-11	1.5E-11	1.0E-11	8.4E-12	7.6E-12	1.4E-12	1.4E-12
NE	4.5E-11	2.5E-11	1.8E-11	1.5E-11	1.4E-11	1.8E-12	1.7E-12
NNE	7.1E-11	3.7E-11	2.6E-11	2.1E-11	1.9E-11	1.9E-12	1.8E-12

Direction	Distance (m)
	1246
N	1.3E-12
NNW	1.3E-12
NW	1.3E-12
WNW	1.3E-12
W	1.3E-12
WSW	1.3E-12
SW	1.3E-12
SSW	1.3E-12
S	1.3E-12
SSE	1.3E-12
SE	1.3E-12
ESE	1.3E-12
E	1.3E-12

		F.sum
ENE	1.3E-12	
NE	1.3E-12	
NNE	1.3E-12	

APPENDIX B

DOE SUPPLEMENTAL INFORMATION

B. DOE SUPPLEMENTAL INFORMATION

- **Provide an estimate of collective dose equivalent for 2002 releases.**
 - The collective population dose for calendar year 2002 emissions to the population within a 10-km distance of the facility is estimated to be 9.7×10^{-5} person-rem/yr.
- **Provide information on the status of compliance with Subparts Q and T of 40 CFR Part 61.**
 - LEHR is in compliance with Subparts Q and T of 40 CFR part 61, based on a radon study conducted at the LEHR Facility by DOE (DOE, 1990).
- **Provide information on radon-220 emissions from sources containing uranium-232 and thorium-232 where emissions potentially can exceed 0.1 mrem/yr to the public or 10% of the nonradon dose to the public.**
 - There are no unencapsulated uranium-232 or thorium-232 sources stored at the facility. The emissions from radon-220 from encapsulated sources would not result in a dose to a member of the public in excess of 0.1 mrem/yr or exceed 10% of the nonradon dose to the public from the Site.
- **Provide information on radon-222 emissions from nondisposal/nonstorage sources where emissions potentially can exceed 0.1 mrem/yr to the public or 10% of the nonradon dose to the public.**
 - There are no nondisposal or nonstorage sources of radon-222 located at the facility.
- **Give the number of emission points subject to the continuous monitoring requirements of Section 61.93(b) of 40 CFR, the number of these emission points that do not comply with Section 61.93(b) requirements, and the cost of upgrades. Describe site periodic confirmatory measurement plans. Indicate the status of the QA program described by Appendix B, Method 114.**
 - There are no point source emissions that require continuous monitoring according to Subpart H of 40 CFR.